



GAZYVA for Previously Untreated Follicular Lymphoma

GAZYVA is a prescription medicine used in combination with chemotherapy, followed by GAZYVA alone in those who responded, to treat stage II bulky, III, or IV follicular lymphoma (FL) in adults who have not had previous FL treatment.

WHAT IS THE MOST IMPORTANT SAFETY INFORMATION I SHOULD KNOW ABOUT GAZYVA?

Tell your doctor right away about any side effect you experience. GAZYVA can cause side effects that can become serious or life-threatening, including:

- **Hepatitis B Virus (HBV):** Hepatitis B can cause liver failure and death. If you have a history of hepatitis B infection, GAZYVA could cause it to return. You should not receive GAZYVA if you have active hepatitis B liver disease. Your doctor or healthcare team will need to screen you for hepatitis B before, and monitor you during and after, your treatment with GAZYVA. Sometimes this will require treatment for hepatitis B. Symptoms of hepatitis include: worsening of fatigue and yellow discoloration of skin or eyes
- **Progressive Multifocal Leukoencephalopathy (PML):** PML is a rare and serious brain infection caused by a virus. PML can be fatal. Your weakened immune system could put you at risk. Your doctor will watch for symptoms. Symptoms of PML include: confusion, difficulty talking or walking, dizziness or loss of balance, and vision problems

*Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including **BOXED WARNINGS**.*


GAZYVA[®]
obinutuzumab
injection | 1,000mg/40mL

What's in this brochure?

This brochure is for adults with advanced* follicular lymphoma, their loved ones, and caregivers.

Inside, you can learn more about this condition, including the symptoms associated with it, common medical tests for diagnosis, the stages of follicular lymphoma, and its treatment goals.

You can also learn about treatment with GAZYVA® (obinutuzumab), how it works, what you can expect from treatment, and about the support and resources that can help you throughout your treatment journey.

Please contact your healthcare treatment team with any questions that you may have.

The information provided in this brochure is meant for informational purposes only. It is not meant to replace your physician's medical advice.

Who should not receive GAZYVA?

Do NOT receive GAZYVA if you have had an allergic reaction (e.g., anaphylaxis or serum sickness) to GAZYVA. Tell your healthcare provider if you have had an allergic reaction to obinutuzumab or any other ingredients in GAZYVA in the past

*"Advanced" means stage II bulky, III, or IV follicular lymphoma.

Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including BOXED WARNINGS.

In this brochure:

About follicular lymphoma	4
Treatment goals and options	10
About GAZYVA	14
What to expect with GAZYVA treatment	18
GAZYVA safety information	22
Glossary	28
Support and resources	30
Talking with your doctor	32
Contacts	33

Underlined, bold medical terms used in this guide will be defined in the Glossary on page 28.


What is follicular lymphoma?

Follicular lymphoma is a cancer of the immune system.

Follicular lymphoma is a type of non-Hodgkin lymphoma—one of the most common forms of blood cancer. Non-Hodgkin lymphoma can be either fast growing (aggressive) or slow growing (indolent).

Follicular lymphoma is the most common slow-growing type and occurs in **lymphocytes**, a type of white blood cell. These cells are an important part of your **immune system** and help defend your body from infection.

When you have follicular lymphoma, too many white blood cells that are affected by the disease build up in your **lymph nodes**, blood, and **bone marrow**. They may also build up in your **spleen** (an organ in your immune system) and cause swelling of this organ.



What are the signs and symptoms of follicular lymphoma?

Although individuals experience symptoms differently, those with follicular lymphoma often will notice an area of painless swelling on the body, such as on the neck, underarm, or groin. Since follicular lymphoma is slow-growing, immediate treatment is not always needed. If you don't appear to have symptoms, you may not be treated right away. This approach is called "active surveillance" or "watchful waiting." When your doctor chooses this approach, he or she will closely monitor your condition.

You can take an active role in keeping an eye on your lymphoma by paying careful attention to changes to your health and communicating symptoms to your doctor.

Possible symptoms you should watch for include (but are not limited to):

- Swollen lymph nodes in the neck, armpit, or groin, which may be painless
- Fever, fatigue, and night sweats
- Unexplained weight loss
- Severe or frequent infections
- Easy bruising or bleeding
- Numbness or tingling in feet and/or hands
- Headaches and/or blurry vision

Your doctor may look for:

- Swelling in your lymph nodes, liver, or spleen
- An increase in the number of abnormal white blood cells
- A decrease in the number of normal blood cells

Symptoms of follicular lymphoma may be seen in other conditions as well. Only your doctor will be able to tell if your symptoms are related to follicular lymphoma.



Understanding medical tests for follicular lymphoma

Follicular lymphoma cannot be diagnosed by symptoms alone. You are not often the first to notice your follicular lymphoma. It is usually detected by routine check-ups or blood work for other health issues.

If your doctor suspects follicular lymphoma after reviewing your signs and symptoms, he or she will order tests to confirm the diagnosis and to determine how advanced the disease is.


Common tests include:



- Routine tests such as physical exam, blood cell counts, and lymph node or bone marrow biopsy
- **Flow cytometry**—a sample of your cells is examined using a laser beam and a computer to find out the type of cancer and the number of cells involved
- Imaging tests such as **PET-CT** or **CT scans**—these scans produce images of the inside of your body that help show the location of follicular lymphoma


The 4 stages of follicular lymphoma

Staging describes how far lymphoma has spread. Keep in mind that even the most advanced stages of follicular lymphoma (stage III and stage IV) are common and can be treated.




STAGE I

One group of lymph nodes is affected either above or below the **diaphragm**.




STAGE II

Two or more groups of lymph nodes are affected either above or below the diaphragm.



STAGE III

Lymph nodes are affected both above and below the diaphragm, or above the diaphragm with spleen involvement.



STAGE IV

Lymphoma is found in bone marrow and/or organs outside of the lymph nodes and spleen.

 **Diaphragm**

*Some patients may be diagnosed with stage II bulky disease. This means that the area of disease is 7 centimeters (about 2.75 inches) or larger.

Goals of treatment for follicular lymphoma

Follicular lymphoma is not curable, but the disease can be managed over time. After initial treatment, your disease may progress (**relapse**) and require treatment again. In each stage of treatment, the goal is to relieve symptoms, stop the cancer from becoming worse, or achieve **remission**.

Treatment goals and options depend on how much your symptoms are affecting you

- **When you have no symptoms**, close monitoring is usually preferred over treatment. This is often referred to as watchful waiting or active surveillance (see page 6)
- **When symptoms appear or worsen**, treatment aims to stop the follicular lymphoma from progressing

Follicular lymphoma can be treated in a number of ways, including antibody therapy and/or chemotherapy. Often, doctors will combine the 2 types of treatment.

Once you begin treatment, your doctor will need to regularly check:



- Your symptoms
- The size of your lymph nodes, liver, or spleen
- Your blood count measures

Treatment options

Treatments may vary from patient to patient and depend on various factors, including the stage of your disease, your age, and overall health. Although patients may respond initially, as many as 1 in 5 patients will relapse within 2 years of receiving their initial treatment therapy. Patients who experience such early relapse have poorer outcomes.

Even after years in remission, follicular lymphoma can recur. When a relapse occurs, the cancer can become more difficult to treat. This means that preventing a patient's disease from returning for as long as possible while minimizing disease symptoms is an important goal of initial treatment.

Please contact your healthcare treatment team with any questions that you may have.



Treatment options (cont'd)

Antibody therapy



Antibody therapy is used to find and destroy specific cells within the body. It can use your body's immune system to help fight cancer. However, it can also harm healthy cells in the body.

Antibody therapy is often an important part of follicular lymphoma treatment plans and can be given along with **chemotherapy**.

How it works

Antibody therapy targets a protein found on the surface of follicular lymphoma cells and certain healthy blood cells. It is thought to use your body's immune system to find and kill lymphoma and healthy cells.

Traditional chemotherapy



Chemotherapy is a type of drug treatment that destroys growing cells, including cancer cells. It is also an important part of follicular lymphoma treatment plans.

Chemotherapy can be given as a single drug or as a combination of drugs, including antibody therapy.

How it works

Chemotherapy attacks growing cancer cells in the body—but also attacks healthy cells, such as those for hair and the intestinal lining.



What is GAZYVA® (obinutuzumab)?

GAZYVA is a type of antibody therapy that targets and attaches to the CD20 proteins found on follicular lymphoma cells as well as some healthy blood cells.

In people with follicular lymphoma, CD20 may be found in higher than normal amounts.

Once attached to the CD20 protein, GAZYVA is thought to work in different ways, including:

- By helping your own immune system destroy the cancer cells
- By destroying the cancer cells directly

GAZYVA can also harm healthy cells in your body.

GAZYVA is a prescription medicine used in combination with chemotherapy, followed by GAZYVA alone in those who responded, to treat stage II bulky, III, or IV follicular lymphoma (FL) in adults who have not had previous FL treatment.

Starting treatment with GAZYVA

On the days you receive GAZYVA, you should expect to spend most of the day at the clinic or infusion center.

Your doctor may have you take medications one hour before you begin taking GAZYVA. Your doctor may refer to this kind of medication as premedication.

Common premedications include acetaminophen, antihistamines, and steroid medications. Taking the suggested medication before treatment may reduce the chance of possible side effects during your GAZYVA infusion. Be sure to check with your medical team about the proper premedication you should take before receiving GAZYVA.



About GAZYVA

How and when am I given GAZYVA® (obinutuzumab)?



GAZYVA is given as an infusion, directly into a vein through a needle in your arm. You'll receive treatment with GAZYVA at a clinic or infusion center.

GAZYVA is given in combination with chemotherapy for a total of 6-8 cycles (depending on the chemotherapy). If you achieve a response after the first 6-8 cycles, your doctor may recommend GAZYVA alone once every 2 months for up to 2 years.

Depending on the type of chemotherapy your doctor recommends, you will receive your medication in one of the following cycles, or rounds, of treatment:

- Six 28-day cycles, when combined with bendamustine
- Six 21-day cycles, followed by 2 additional 21-day cycles of GAZYVA alone, when combined with CHOP
- Eight 21-day cycles, when combined with CVP

Treatment occurs only on certain days of each cycle and is usually followed by several weeks of rest and recovery.

During the infusion process, some patients may experience side effects. Please refer to page 20 for more information about these reactions.

Speak with your doctor if you miss a dose of GAZYVA.

Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including BOXED WARNINGS.

The chart below shows when you should receive your GAZYVA infusion

CYCLE 1

WEEK 1	DAY 1 	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
WEEK 2	DAY 8 	DAY 9	DAY 10	DAY 11	DAY 12	DAY 13	DAY 14
WEEK 3	DAY 15 	DAY 16	DAY 17	DAY 18	DAY 19	DAY 20	DAY 21
WEEK 4	DAY 22	DAY 23	DAY 24	DAY 25	DAY 26	DAY 27	DAY 28

For Cycle 1, you will receive GAZYVA on Days 1, 8, and 15

CYCLES 2-6 or 2-8

WEEK 1	DAY 1 	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6	DAY 7
WEEK 2	DAY 8	DAY 9	DAY 10	DAY 11	DAY 12	DAY 13	DAY 14
WEEK 3	DAY 15	DAY 16	DAY 17	DAY 18	DAY 19	DAY 20	DAY 21
WEEK 4	DAY 22	DAY 23	DAY 24	DAY 25	DAY 26	DAY 27	DAY 28

For Cycles 2-6 or 2-8, depending on the type of chemotherapy your doctor recommends, you will receive GAZYVA on the first day of each cycle

If you complete your GAZYVA cycles in combination with chemotherapy and your disease has responded to treatment, your doctor may recommend GAZYVA once every 2 months for up to 2 years

GAZYVA 1,000 mg

Preparing for your infusion

One week before:

- Tell your doctor about all the medicines you take, including prescription and non-prescription medicine, vitamins, and herbal supplements
- **Speak to your doctor if you take medications to control your blood pressure. Your doctor may instruct you not to take them on the day of your GAZYVA infusion**
- Arrange a ride. After your infusion, you may feel drowsy or dizzy. It is a good idea to have someone else drive you home

One day before:

- Remember to follow any changes your doctor has made to your regular medication schedule

The day of your infusion



At home:

- Wear or bring loose clothing so you can be comfortable during your infusion
- Pack food and drink—bringing some snacks or a packed meal can help you get through the day
- Bring a book or activities to help pass the time



At the clinic or infusion center:

- Take the medications your doctor prescribed to help reduce the side effects of the infusion process
- Relax during your infusion, but be aware of what is happening to your body. Tell your doctor or nurse right away if you feel any side effects, including the symptoms of an infusion-related reaction



Preparing for your infusion

Infusion-Related Reactions (IRRs)

These side effects may occur during or within 24 hours of any GAZYVA infusion. Some IRRs can be serious. These may include, but are not limited to, severe allergic reactions (anaphylaxis), acute life-threatening breathing problems, or other life-threatening IRRs.

If you have a reaction, your infusion will either be slowed or stopped until your symptoms are resolved. Most patients are able to complete infusions and receive medication again. However, if the infusion-related reaction is serious, the infusion of GAZYVA will be permanently stopped.

Your healthcare team will take steps to help lessen any side effects that you may have to the infusion process. You may be given medicines to take before each GAZYVA treatment.

Symptoms of IRRs may include:



- Fast heartbeat
- Tiredness
- Dizziness
- Headache
- Redness of the face
- Nausea
- Chills
- Fever
- Vomiting
- Diarrhea
- Rash
- High blood pressure
- Low blood pressure
- Difficulty breathing
- Chest discomfort



Important Safety Information about GAZYVA® (obinutuzumab)

What is the most important safety information I should know about GAZYVA?

Tell your doctor right away about any side effect you experience. GAZYVA can cause side effects that can become serious or life threatening, including:

- **Hepatitis B Virus (HBV):** Hepatitis B can cause liver failure and death. If you have a history of hepatitis B infection, GAZYVA could cause it to return. You should not receive GAZYVA if you have active hepatitis B liver disease. Your doctor or healthcare team will need to screen you for hepatitis B before, and monitor you during and after, your treatment with GAZYVA. Sometimes this will require treatment for hepatitis B. Symptoms of hepatitis include: worsening of fatigue and yellow discoloration of skin or eyes
- **Progressive Multifocal Leukoencephalopathy (PML):** PML is a rare and serious brain infection caused by a virus. PML can be fatal. Your weakened immune system could put you at risk. Your doctor will watch for symptoms. Symptoms of PML include: confusion, difficulty talking or walking, dizziness or loss of balance, and vision problems

Who should not receive GAZYVA?

- **Do NOT** receive GAZYVA if you have had an allergic reaction (e.g., anaphylaxis or serum sickness) to GAZYVA. Tell your healthcare provider if you have had an allergic reaction to obinutuzumab or any other ingredients in GAZYVA in the past

Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including BOXED WARNINGS.

What are additional possible serious side effects of GAZYVA?

Tell your doctor right away about any side effect you experience. GAZYVA can cause side effects that may become severe or life threatening, including:

- **Infusion-Related Reactions (IRRs):** These side effects may occur during or within 24 hours of any GAZYVA infusion. Some IRRs can be serious, including, but not limited to, severe allergic reactions (anaphylaxis), acute life-threatening breathing problems, or other life-threatening IRRs. If you have a reaction, the infusion is either slowed or stopped until your symptoms are resolved. Most patients are able to complete infusions and receive medication again. However, if the infusion-related reaction is life threatening, the infusion of GAZYVA will be permanently stopped. Your healthcare team will take steps to help lessen any side effects you may have to the infusion process. You may be given medicines to take before each GAZYVA treatment. Symptoms of IRRs may include: fast heartbeat, tiredness, dizziness, headache, redness of the face, nausea, chills, fever, vomiting, constipation, diarrhea, rash, high blood pressure, low blood pressure, difficulty breathing, and chest discomfort
- **Hypersensitivity Reactions Including Serum Sickness:** Some people receiving GAZYVA may have severe or life-threatening allergic reactions. This reaction may be severe, may happen during or after an infusion, and may affect many areas of the body. If an allergic reaction occurs, your doctor will stop the infusion and permanently discontinue GAZYVA
- **Tumor Lysis Syndrome (TLS):** Tumor lysis syndrome, including fatal cases, has been reported in patients receiving GAZYVA. GAZYVA works to break down cancer cells quickly. As cancer cells break apart, their contents are released into the blood. These contents may cause damage to organs and the heart, and may lead to kidney failure requiring the need for dialysis treatment. Your doctor may prescribe medication to help prevent TLS. Your doctor will also conduct regular blood tests to check for TLS. Symptoms of TLS may include nausea, vomiting, diarrhea, and tiredness

Important Safety Information about GAZYVA® (obinutuzumab) (cont'd)

- **Infections:** While you're taking GAZYVA, you may develop infections. Some of these infections can be fatal and severe, so be sure to talk to your doctor if you think you have an infection. Patients administered GAZYVA in combination with chemotherapy, followed by GAZYVA alone are at a high risk of infections during and after treatment. Patients with a history of recurring or chronic infections may be at an increased risk of infection. Patients with an active infection should not be treated with GAZYVA. Patients taking GAZYVA plus bendamustine may be at a higher risk for fatal or severe infections compared to patients taking GAZYVA plus CHOP or CVP
- **Low White Blood Cell Count:** When you have an abnormally low count of infection-fighting white blood cells, it is called neutropenia. While you are taking GAZYVA, your doctor will do blood work to check your white blood cell count. Severe and life-threatening neutropenia can develop during or after treatment with GAZYVA. Some cases of neutropenia can last for more than one month. If your white blood cell count is low, your doctor may prescribe medication to help prevent infections

- **Low Platelet Count:** Platelets help stop bleeding or blood loss. GAZYVA may reduce the number of platelets you have in your blood; having a low platelet count is called thrombocytopenia. This may affect the clotting process. While you are taking GAZYVA, your doctor will do blood work to check your platelet count. Severe and life-threatening thrombocytopenia can develop during treatment with GAZYVA. Fatal bleeding events have occurred in patients treated with GAZYVA. If your platelet count gets too low, your treatment may be delayed or reduced

The most common side effects seen with GAZYVA in a study that included previously untreated FL patients were infusion-related reactions, low white blood cell count, upper respiratory tract infections, cough, constipation and diarrhea

What other information should I tell my doctor before receiving GAZYVA?

You should talk to your doctor about:

- **Immunizations:** Before receiving GAZYVA therapy, tell your healthcare provider if you have recently received or are scheduled to receive a vaccine. People who are treated with GAZYVA should not receive live vaccines
- **Pregnancy:** Tell your doctor if you are pregnant, think that you might be pregnant, or plan to become pregnant, or are breastfeeding. GAZYVA may harm your unborn baby. Speak to your doctor about using GAZYVA while you are pregnant. Talk to your doctor or your child's doctor about the safety and timing of live virus vaccinations to your infant if you received GAZYVA during pregnancy. It is not known if GAZYVA may pass into your breast milk. Women of childbearing potential should use effective contraception while taking GAZYVA and for 6 months after your GAZYVA treatment
- **Breastfeeding:** Because of the potential risk of serious side reactions in breastfed children, women should not breastfeed while taking GAZYVA and for 6 months after your last dose

Important Safety Information about GAZYVA[®] (obinutuzumab) (cont'd)

Tell your doctor about any side effects.

These are not all of the possible side effects of GAZYVA. For more information, ask your doctor or pharmacist.

GAZYVA is available by prescription only.

You may report side effects to the FDA at (800) FDA-1088, or www.fda.gov/medwatch.

You may also report side effects to Genentech at (888) 835-2555.

Please see the accompanying full Prescribing Information, including BOXED WARNINGS, for additional Important Safety Information.

Remember to tell your healthcare team exactly how you feel. This will allow them to take the proper actions to help you.

Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including BOXED WARNINGS.



Important terms found in this guide.

Antibody therapy: A type of therapy used to find and destroy specific cells within the body (for example, the cells where follicular lymphoma starts). Antibody therapy can also harm healthy cells in the body.

Bone marrow: The soft spongy material that fills the inside of bones. Bone marrow is the source of new blood cells, and platelets are made in the bone marrow.

Chemotherapy: A drug treatment that destroys growing cells, including cancer cells.

CT (computed tomography) scan: A medical test that uses a computer linked to an x-ray machine to take pictures of the inside of the body.

Diaphragm: The muscle that divides the chest cavity from the abdominal cavity.

Flow cytometry: This test identifies the type of blood cancer and number of cells involved.

Follicular lymphoma: The most common type of indolent (slow growing) non-Hodgkin's lymphoma (NHL). With NHL, abnormal lymphocytes build up in the lymph nodes, bone marrow, and spleen. Follicular lymphoma can be a chronic disease.

Immune system: The group of organs and cells that defends the body from infections.

Lymph node: A small bean-shaped organ that stores white blood cells.

Lymphocytes: A type of white blood cell that plays an important role in fighting infection.

PET (Positron emission tomography) scan: A medical test that uses a form of radioactive sugar that can be traced by a special camera. This can help tell if an area contains lymphoma and if that lymphoma is responding to treatment.

PET-CT (Positron emission tomography-computed tomography) scan: A medical test that combines the PET scan with a CT scan. This lets the doctor compare areas with lymphoma on the PET scan with the more detailed appearance of that area on the CT scan.

Relapse: The return of a disease, or the signs and symptoms of a disease after a period of improvement.

Remission: A term used to describe a response to treatment. Partial remission means the cancer is significantly improved, but evidence of the cancer remains. Complete remission means all evidence of the cancer is gone for a period of time.

Spleen: An organ that is part of the lymphatic system. The spleen makes lymphocytes, filters blood, stores blood cells, and destroys old blood cells. It is located on the left side of the abdomen near the stomach.



Support from Genentech



Genentech Oncology Co-pay Assistance Program*

Genentech co-pay programs help people with commercial health insurance. This might be a plan you get through your employer or one you purchased through a Health Insurance Marketplace like HealthCare.gov. To qualify, you must also meet other criteria.

*This Co-pay Program is valid ONLY for patients with commercial insurance who have a valid prescription for a Food and Drug Administration (FDA)-approved indication of a Genentech medication. Patients using Medicare, Medicaid, or any other federal or state government program to pay for their medications are not eligible. Under the Program, the patient will pay a co-pay. After reaching the maximum Program benefit, the patient will be responsible for all out-of-pocket costs. All participants are responsible for reporting the receipt of all Program benefits as required by any insurer or by law. No party may seek reimbursement for all or any part of the benefit received through this Program. This Program is void where prohibited by law. Genentech reserves the right to rescind, revoke, or amend the Program without notice at any time. Additional eligibility criteria apply. See full terms and conditions at copayassistancenow.com.

Independent Co-pay Assistance Foundations†

If you need help with the co-pay for your Genentech medicine, Genentech Access Solutions can refer you to an independent co-pay assistance foundation. Independent co-pay assistance foundations help patients with public or commercial health insurance.

†Independent co-pay assistance foundations have their own rules for eligibility. We cannot guarantee a foundation will help you. We only can refer you to a foundation that supports your disease state. We do not endorse or show financial preference for any particular foundation. The foundations we refer you to are not the only ones that might be able to help you.

Genentech Patient Foundation‡

The Genentech Patient Foundation gives free Genentech medicine to people who need it.

‡If you have health insurance, you must have already tried other types of financial assistance. You also need to meet income requirements. If you do not have insurance, or if your insurance does not cover your Genentech medicine, you must meet different income requirements.

To learn more about how we can help, **CALL (877) GENENTECH/(877) 436-3683**
VISIT www.genentech-access.com/GAZYVA/patients

Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including BOXED WARNINGS.

Helpful resources

Many patient support groups offer helpful information about cancer. Some may also help you connect with a local support group. You can share your experiences and learn more about follicular lymphoma. Many people find this helps them stay informed and stay positive.

Cancer organizations

American Cancer Society

1-800-ACS-2345 (1-800-227-2345)
www.cancer.org

CancerCare, Inc.

1-800-813-HOPE (1-800-813-4673)
www.cancer.org

National Cancer Institute

1-800-4-CANCER (1-800-422-6237)
www.cancer.gov

National Comprehensive Cancer Network

www.nccn.org/patients

Genentech does not control or endorse third-party organizations. The information provided by Genentech or these organizations is meant for informational purposes only. It is not meant to replace your doctor's medical advice.

Lymphoma organizations

Lymphoma Research Foundation

1-800-500-9976
www.lymphoma.org

The Leukemia & Lymphoma Society

1-800-955-4572
www.lls.org

Support organizations

Cancer Hope Network

1-877-HOPENET (1-877-467-3638)
www.cancerhopenetwork.org

Cancer Support Community

1-888-793-WELL (1-888-793-9355)
www.cancersupportcommunity.org

Patient Advocate Foundation

1-800-532-5274
www.patientadvocate.org

Questions to ask your doctor



It is important to ask your doctor or nurse any questions you may have about your condition and treatment. It's a good idea to write down a list of questions before your appointment. It also may be a good idea to bring a family member or friend to your appointment for support—and to help you keep track of the answers.

Here are some questions to consider asking:

- What are my treatment options?
- What will my treatment schedule be?
- Do I need to take medication at home as well?
- What are the possible side effects of treatment?
- Is there more I can do to make the most of my treatment?
- Where can I get more information about previously untreated follicular lymphoma?
- Where can I find information about support to help me pay for my GAZYVA® (obinutuzumab) therapy?
- How will treatment with GAZYVA affect my lifestyle (working, traveling, etc)?

Use this space to help keep track of important contact information

Doctor (Name): _____



Nurse (Name): _____



Nurse (Name): _____



Hospital (Name): _____



Pharmacy (Name): _____



Emergency (Name): _____





GAZYVA[®]
obinutuzumab
injection | 1,000mg/40mL

For more information, visit www.GAZYVA.com

Images in this brochure do not depict actual patients or healthcare providers.

Please see additional Important Safety Information throughout as well as accompanying full Prescribing Information, including BOXED WARNINGS.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use GAZYVA safely and effectively. See full prescribing information for GAZYVA.

GAZYVA® (obinutuzumab) injection, for intravenous use
Initial U.S. Approval: 2013

WARNING: HEPATITIS B VIRUS REACTIVATION and PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY
See full prescribing information for complete boxed warning.

- **Hepatitis B Virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death. (5.1)**
- **Progressive Multifocal Leukoencephalopathy (PML) resulting in death. (5.2)**

RECENT MAJOR CHANGES

Warnings and Precautions, Embryo-Fetal Toxicity (5.10)-----03/2020

INDICATIONS AND USAGE

GAZYVA (obinutuzumab) is a CD20-directed cytolytic antibody indicated:

- in combination with chlorambucil, for the treatment of patients with previously untreated chronic lymphocytic leukemia. (1, 14)
- in combination with bendamustine followed by GAZYVA monotherapy, for the treatment of patients with follicular lymphoma who relapsed after, or are refractory to, a rituximab-containing regimen. (1, 14)
- in combination with chemotherapy followed by GAZYVA monotherapy in patients achieving at least a partial remission, for the treatment of adult patients with previously untreated stage II bulky, III or IV follicular lymphoma. (1, 14)

DOSAGE AND ADMINISTRATION

- Premedicate for infusion-related reactions and tumor lysis syndrome. (2.1, 5.3, 5.4)
- Administer only as intravenous infusion. Do not administer as an intravenous push or bolus. (2.1)
- The recommended dosage for chronic lymphocytic leukemia is 100 mg on day 1 and 900 mg on day 2 of Cycle 1, 1,000 mg on day 8 and 15 of Cycle 1, and 1,000 mg on day 1 of Cycles 2–6. (2.2)
- The recommended dosage for follicular lymphoma is 1,000 mg on day 1, 8 and 15 of Cycle 1, 1,000 mg on day 1 of Cycles 2-6 or Cycles 2-8, and then 1,000 mg every 2 months for up to 2 years. (2.3)

DOSAGE FORMS AND STRENGTHS

Injection: 1,000 mg/40 mL (25 mg/mL) single-dose vial. (3)

CONTRAINDICATIONS

GAZYVA is contraindicated in patients with known hypersensitivity reactions (e.g., anaphylaxis) to obinutuzumab or any of the excipients, including serum sickness with prior obinutuzumab use. (4)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: HEPATITIS B VIRUS REACTIVATION AND PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY

1 INDICATIONS AND USAGE

- 1.1 Chronic Lymphocytic Leukemia
- 1.2 Follicular Lymphoma

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosing Information
- 2.2 Recommended Dosage for Chronic Lymphocytic Leukemia
- 2.3 Recommended Dosage for Follicular Lymphoma
- 2.4 Recommended Premedication and Prophylactic Medications
- 2.5 Dosage Modifications for Adverse Reactions
- 2.6 Preparation and Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hepatitis B Virus Reactivation
- 5.2 Progressive Multifocal Leukoencephalopathy
- 5.3 Infusion-Related Reactions
- 5.4 Hypersensitivity Reactions Including Serum Sickness
- 5.5 Tumor Lysis Syndrome
- 5.6 Infections
- 5.7 Neutropenia
- 5.8 Thrombocytopenia
- 5.9 Immunizations
- 5.10 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

WARNINGS AND PRECAUTIONS

- **Infusion-Related Reactions:** Premedicate patients with glucocorticoid, acetaminophen, and anti-histamine. Monitor patients closely during infusions. Interrupt, reduce rate, or discontinue for infusion-related reactions based on severity. (2.1, 5.3)
- **Hypersensitivity Reactions Including Serum Sickness:** Discontinue GAZYVA permanently. (5.4)
- **Tumor Lysis Syndrome:** Premedicate with anti-hyperuricemics and adequate hydration, especially for patients with high tumor burden, high circulating lymphocyte count or renal impairment. Correct electrolyte abnormalities, provide supportive care, and monitor renal function and fluid balance. (5.5)
- **Infections:** Do not administer GAZYVA to patients with an active infection. Patients with a history of recurring or chronic infections may be at increased risk of infection. (5.6)
- **Neutropenia:** In patients with Grade 3 to 4 neutropenia, monitor laboratory tests until resolution and for infection. Consider dose delays and infection prophylaxis, as appropriate. (5.7)
- **Thrombocytopenia:** Monitor platelet counts and for bleeding. Transfusion may be necessary. (5.8)
- **Immunization:** Avoid administration of live virus vaccines during GAZYVA treatment and until B-cell recovery. (5.9)
- **Embryo-Fetal Toxicity:** Can cause fetal harm. Advise females of reproductive potential of the potential risk to a fetus and use effective contraception. (5.10)

ADVERSE REACTIONS

The most common adverse reactions (incidence $\geq 20\%$ and $\geq 2\%$ greater in the GAZYVA treated arm) were:

- **Previously untreated CLL:** infusion-related reactions and neutropenia. (6)
- **Relapsed or refractory NHL:** infusion-related reactions, fatigue, neutropenia, cough, upper respiratory tract infections, and musculoskeletal pain. (6)
- **Previously untreated NHL:** infusion-related reactions, neutropenia, upper respiratory tract infections, cough, constipation, and diarrhea. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Genentech at 1-888-835-2555 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- **Lactation:** Advise not to breastfeed. (8.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 3/2020

6.1 Clinical Trial Experience

6.2 Immunogenicity

6.3 Additional Clinical Trial Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Chronic Lymphocytic Leukemia

14.2 Follicular Lymphoma

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied/Storage

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: HEPATITIS B VIRUS REACTIVATION and PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY

- **Hepatitis B Virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients receiving CD20-directed cytolytic antibodies, including GAZYVA. Screen all patients for HBV infection before treatment initiation. Monitor HBV-positive patients during and after treatment with GAZYVA. Discontinue GAZYVA and concomitant medications in the event of HBV reactivation [see Warnings and Precautions (5.1)].**
- **Progressive Multifocal Leukoencephalopathy (PML) including fatal PML, can occur in patients receiving GAZYVA [see Warnings and Precautions (5.2)].**

1 INDICATIONS AND USAGE

1.1 Chronic Lymphocytic Leukemia (CLL)

GAZYVA, in combination with chlorambucil, is indicated for the treatment of patients with previously untreated chronic lymphocytic leukemia.

1.2 Follicular Lymphoma (FL)

GAZYVA, in combination with bendamustine followed by GAZYVA monotherapy, is indicated for the treatment of patients with follicular lymphoma who relapsed after, or are refractory to, a rituximab-containing regimen.

GAZYVA, in combination with chemotherapy followed by GAZYVA monotherapy in patients achieving at least a partial remission, is indicated for the treatment of adult patients with previously untreated stage II bulky, III or IV follicular lymphoma.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosing Information

- Premedicate before each infusion [see *Dosage and Administration (2.4)*].
- Provide prophylactic hydration and anti-hyperuricemics to patients at high risk of tumor lysis syndrome [see *Dosage and Administration (2.4)* and *Warnings and Precautions (5.4)*].
- Administer only as an intravenous infusion through a dedicated line [see *Dosage and Administration (2.6)*].
- Do not administer as an intravenous push or bolus.
- Monitor blood counts at regular intervals.
- GAZYVA should only be administered by a healthcare professional with appropriate medical support to manage severe infusion-related reactions that can be fatal if they occur [see *Warnings and Precautions (5.3)*].

2.2 Recommended Dosage for Chronic Lymphocytic Leukemia

Each dose of GAZYVA is 1,000 mg administered intravenously with the exception of the first infusions in Cycle 1, which are administered on day 1 (100 mg) and day 2 (900 mg) according to Table 1.

Table 1 Dose of GAZYVA to be Administered During Six 28-Day Treatment Cycles for Patients with CLL

Day of treatment cycle		Dose of GAZYVA	Rate of infusion
Cycle 1 (loading doses)	Day 1	100 mg	Administer at 25 mg/hr over 4 hours. Do not increase the infusion rate.
	Day 2	900 mg	If no infusion-related reaction occurred during the previous infusion, administer at 50 mg/hr. The rate of the infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr. If an infusion-related reaction occurred during the previous infusion, administer at 25 mg/hr. The rate of infusion can be escalated in increments of up to 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.
	Day 8	1,000 mg	If no infusion-related reaction occurred during the previous infusion and the final infusion rate was 100 mg/hr or faster, infusions can be started at a rate of 100 mg/hr and increased by 100 mg/hr increments every 30 minutes to a maximum of 400 mg/hr. If an infusion-related reaction occurred during the previous infusion, administer at 50 mg/hr. The rate of infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.
	Day 15	1,000 mg	
Cycles 2–6	Day 1	1,000 mg	If no infusion-related reaction occurred during the previous infusion, administer at 50 mg/hr. The rate of infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.

If a planned dose of GAZYVA is missed, administer the missed dose as soon as possible and adjust dosing schedule to maintain the time interval between doses. If appropriate, patients who do not complete the Day 1 Cycle 1 dose may proceed to the Day 2 Cycle 1 dose.

2.3 Recommended Dosage for Follicular Lymphoma

Each dose of GAZYVA is 1,000 mg administered intravenously according to Table 2.

For patients with relapsed or refractory FL, administer GAZYVA in combination with bendamustine in six 28-day cycles. Patients who achieve stable disease, complete response, or partial response to the initial 6 cycles should continue on GAZYVA 1,000 mg as monotherapy for up to two years.

For patients with previously untreated FL, administer GAZYVA with one of the following chemotherapy regimens:

- Six 28-day cycles in combination with bendamustine
- Six 21-day cycles in combination with CHOP, followed by 2 additional 21-day cycles of GAZYVA alone
- Eight 21-day cycles in combination with CVP

Patients with previously untreated FL who achieve a complete response or partial response to the initial 6 or 8 cycles should continue on GAZYVA 1,000 mg as monotherapy for up to two years.

Table 2 Dose of GAZYVA to be Administered During 6–8 Treatment Cycles, Followed by GAZYVA Monotherapy for Patients with FL

Day of treatment cycle		Dose of GAZYVA	Rate of infusion
Cycle 1 (loading doses)	Day 1	1,000 mg	Administer at 50 mg/hr. The rate of the infusion can be escalated in 50 mg/hr increments every 30 minutes to a maximum of 400 mg/hr.
	Day 8	1,000 mg	If no infusion-related reaction or an infusion-related reaction of Grade 1 occurred during the previous infusion and the final infusion rate was 100 mg/hr or faster, infusions can be started at a rate of 100 mg/hr and increased by 100 mg/hr increments every 30 minutes to a maximum of 400 mg/hr.
	Day 15	1,000 mg	
Cycles 2–6 or 2–8	Day 1	1,000 mg	If an infusion-related reaction of Grade 2 or higher occurred during the previous infusion, administer at 50 mg/hr. The rate of infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.
Monotherapy	Every two months for up to two years	1,000 mg	

If a planned dose of GAZYVA is missed, administer the missed dose as soon as possible. During GAZYVA and chemotherapy treatment, adjust the dosing schedule accordingly to maintain the time interval between chemotherapy cycles. During monotherapy, maintain the original dosing schedule for subsequent doses. Initiate monotherapy approximately two months after the last dose of GAZYVA administered during the induction phase.

2.4 Recommended Premedication and Prophylactic Medications

Infusion-Related Reactions

Premedication to reduce the risk of infusion-related reactions is outlined in Table 3 [see *Warnings and Precautions (5.3)*].

Hypotension may occur during GAZYVA intravenous infusions. Consider withholding antihypertensive treatments for 12 hours prior to and throughout each GAZYVA infusion and for the first hour after administration [see *Warnings and Precautions (5.3)*].

Table 3 Premedication for GAZYVA Infusion to Reduce Infusion-Related Reactions (IRR)

Day of Treatment Cycle	Patients requiring premedication	Premedication	Administration
Cycle 1: CLL Day 1, Day 2 FL Day 1	All patients	Intravenous glucocorticoid: 20 mg dexamethasone or 80 mg methylprednisolone ^{1,2}	Completed at least 1 hour prior to GAZYVA infusion.
		650–1,000 mg acetaminophen	
		anti-histamine (e.g., 50 mg diphenhydramine)	At least 30 minutes before GAZYVA infusion.
All subsequent	All patients	650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.

infusions, CLL or FL	Patients with an IRR (Grade 1-2) with the previous infusion	650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.
		anti-histamine (e.g., 50 mg diphenhydramine)	
	Patients with a Grade 3 IRR with the previous infusion OR with a lymphocyte count > 25 x 10 ⁹ /L prior to next treatment	Intravenous glucocorticoid: 20 mg dexamethasone or 80 mg methylprednisolone ¹	Completed at least 1 hour prior to GAZYVA infusion.
		650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.
anti-histamine (e.g., 50 mg diphenhydramine)			

¹ Hydrocortisone is not recommended as it has not been effective in reducing the rate of infusion-related reactions.

² If a glucocorticoid-containing chemotherapy regimen is administered on the same day as GAZYVA, the glucocorticoid can be administered as an oral medication if given at least 1 hour prior to GAZYVA, in which case additional intravenous glucocorticoid as premedication is not required.

Tumor Lysis Syndrome Prophylaxis

Patients with high tumor burden, high circulating absolute lymphocyte counts (greater than 25 x 10⁹/L) or renal impairment are considered at risk of tumor lysis syndrome and should receive prophylaxis. Premedicate with anti-hyperuricemics (e.g., allopurinol or rasburicase) and ensure adequate hydration prior to start of GAZYVA therapy. Continue prophylaxis prior to each subsequent GAZYVA infusion, as needed [see *Warnings and Precautions (5.4)*].

Antimicrobial Prophylaxis

Patients with Grade 3 to 4 neutropenia lasting more than one week are strongly recommended to receive antimicrobial prophylaxis until resolution of neutropenia to Grade 1 or 2. Consider antiviral and antifungal prophylaxis for patients with severe and long lasting (> 1 week) neutropenia.

2.5 Dosage Modifications for Adverse Reactions

Infusion-Related Reactions

If a patient experiences an infusion-related reaction (IRR), adjust the infusion as follows [see *Warnings and Precautions (5.3)*]:

- Grade 4 (life-threatening): Stop infusion immediately and permanently discontinue GAZYVA.
- Grade 3 (severe): Interrupt infusion and manage symptoms. Upon resolution of symptoms, consider restarting GAZYVA infusion at no more than half the previous rate (the rate being used at the time that the IRR reaction occurred), and if patient does not experience any further IRR symptoms, infusion rate escalation may resume at the increments and intervals as appropriate for the treatment cycle dose. Permanently discontinue treatment if patients experience a Grade 3 IRR at rechallenge.
 - For CLL patients only, the Day 1 infusion rate may be increased back up to 25 mg/hr after 1 hour but not increased further.
- Grade 1–2 (mild to moderate): Reduce infusion rate or interrupt infusion and manage symptoms. Upon resolution of symptoms, continue or resume GAZYVA infusion, and if patient does not experience any further IRR symptoms, infusion rate escalation may resume at the increments and intervals as appropriate for the treatment cycle dose.

- For CLL patients only, the Day 1 infusion rate may be increased back up to 25 mg/hr after 1 hour but not increased further.

Other Adverse Reactions

Consider treatment interruption if patients experience an infection, Grade 3 or 4 cytopenia, or a \geq Grade 2 non-hematologic toxicity.

2.6 Preparation and Administration

Preparation

Prepare the solution for infusion, using aseptic technique, as follows:

- Inspect visually for any particulate matter and discoloration prior to administration.
- Dilute into a 0.9% sodium chloride PVC or non-PVC polyolefin infusion bag.

Chronic Lymphocytic Leukemia

- Preparation of solution for infusion on day 1 (100 mg) and day 2 (900 mg) of Cycle 1:
 - Prepare day 1 (100 mg) and day 2 (900 mg) infusion bags at the same time using one vial (1,000 mg/40 mL) on day 1
 - Withdraw 40 mL of GAZYVA solution from the vial.
 - Dilute 4 mL (100 mg) of GAZYVA into a 100 mL 0.9% Sodium Chloride Injection, USP infusion bag for immediate administration.
 - Dilute the remaining 36 mL (900 mg) into a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag at the same time for use on day 2 and store at 2°C to 8°C (36°F to 46°F) for up to 24 hours. After allowing the diluted bag to come to room temperature, use immediately.
 - Clearly label each infusion bag.
- Preparation of solution for infusion on day 8 and 15 of Cycle 1 and day 1 of Cycles 2–6:
 - Withdraw 40 mL of GAZYVA solution from the vial.
 - Dilute 40 mL (1,000 mg) into a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag.

Follicular Lymphoma

- Preparation of solution for infusion:
 - Withdraw 40 mL of GAZYVA solution from the vial.
 - Dilute 40 mL (1,000 mg) into a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag.
- Mix diluted solution by gentle inversion. Do not shake or freeze.
- For microbiological stability, immediately use diluted GAZYVA infusion solution. If not used immediately, store in a refrigerator at 2°C to 8°C (36°F to 46°F) for up to 24 hours prior to use.

The product can be administered at a final concentration of 0.4 mg/mL to 4 mg/mL.

Storage

Use the diluted solution immediately. If not used immediately, store for up to 24 hours at 2°C to 8°C. Discard after 24 hours.

Administration

- Administer as an intravenous infusion only.
- Do not administer as an intravenous push or bolus.
- Do not mix GAZYVA with other drugs.
- No incompatibilities between GAZYVA and polyvinylchloride (PVC) or non-PVC polyolefin bags and administration sets have been observed.

3 DOSAGE FORMS AND STRENGTHS

Injection: 1,000 mg/40 mL (25 mg/mL) clear, colorless to slightly brown solution in single-dose vial.

4 CONTRAINDICATIONS

GAZYVA is contraindicated in patients with known hypersensitivity reactions (e.g., anaphylaxis) to obinutuzumab or to any of the excipients, or serum sickness with prior obinutuzumab use [*see Warnings and Precautions Section (5.4)*].

5 WARNINGS AND PRECAUTIONS

5.1 Hepatitis B Virus Reactivation

Hepatitis B virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients treated with anti-CD20 antibodies such as GAZYVA. HBV reactivation has been reported in patients who are hepatitis B surface antigen (HBsAg) positive and also in patients who are HBsAg negative but are hepatitis B core antibody (anti-HBc) positive. Reactivation has also occurred in patients who appear to have resolved hepatitis B infection (i.e., HBsAg negative, anti-HBc positive, and hepatitis B surface antibody [anti-HBs] positive).

HBV reactivation is defined as an abrupt increase in HBV replication manifesting as a rapid increase in serum HBV DNA level or detection of HBsAg in a person who was previously HBsAg negative and anti-HBc positive. Reactivation of HBV replication is often followed by hepatitis, i.e., increase in transaminase levels and, in severe cases, increase in bilirubin levels, liver failure, and death.

Screen all patients for HBV infection by measuring HBsAg and anti-HBc before initiating treatment with GAZYVA. For patients who show evidence of hepatitis B infection (HBsAg positive [regardless of antibody status] or HBsAg negative but anti-HBc positive), consult healthcare providers with expertise in managing hepatitis B regarding monitoring and consideration for HBV antiviral therapy.

Monitor patients with evidence of current or prior HBV infection for clinical and laboratory signs of hepatitis or HBV reactivation during and for several months following treatment with GAZYVA. HBV reactivation has been reported for other CD20-directed cytolytic antibodies following completion of therapy.

In patients who develop reactivation of HBV while receiving GAZYVA, immediately discontinue GAZYVA and any concomitant chemotherapy and institute appropriate treatment. Resumption of GAZYVA in patients whose HBV reactivation resolves should be discussed with healthcare providers with expertise in managing hepatitis B. Insufficient data exist regarding the safety of resuming GAZYVA in patients who develop HBV reactivation.

5.2 Progressive Multifocal Leukoencephalopathy

John Cunningham (JC) virus infection resulting in progressive multifocal leukoencephalopathy (PML), which can be fatal, occurred in patients treated with GAZYVA. Consider the diagnosis of PML in any patient presenting with new onset or changes to preexisting neurologic manifestations. Evaluation of PML includes, but is not limited to, consultation with a neurologist, brain MRI, and lumbar puncture. Discontinue GAZYVA therapy and consider discontinuation or reduction of any concomitant chemotherapy or immunosuppressive therapy in patients who develop PML.

5.3 Infusion-Related Reactions

GAZYVA can cause severe and life-threatening infusion-related reactions (IRRs). Sixty-five percent of patients with CLL experienced a reaction to the first 1,000 mg of GAZYVA infused. Thirty-seven percent of patients with relapsed or refractory NHL and 60% of patients with previously untreated NHL experienced a reaction on Day 1 of GAZYVA infusion. IRRs have occurred within 24 hours of receiving GAZYVA. IRRs can also occur with subsequent infusions. Symptoms may include hypotension, tachycardia, dyspnea, and respiratory symptoms (e.g., bronchospasm, larynx and throat irritation, wheezing, laryngeal edema). The most frequently reported symptoms include nausea, fatigue, chest discomfort, dyspnea, dizziness, vomiting, diarrhea, rash, hypertension, hypotension, flushing, headache, pyrexia, and chills [see *Adverse Reactions (6.1)*].

Premedicate patients with acetaminophen, antihistamine, and a glucocorticoid [see *Dosage and Administration (2.4)*]. Closely monitor patients during the entire infusion. Reduce infusion rate, interrupt infusion or permanently discontinue GAZYVA for IRRs based on severity [see *Dosage and Administration (2.5)*]. Institute medical management (e.g., glucocorticoids, epinephrine, bronchodilators, and/or oxygen) for IRRs as needed.

For patients with preexisting cardiac or pulmonary conditions, monitor more frequently throughout the infusion and the post-infusion period since they may be at greater risk of experiencing more severe reactions. Hypotension may occur as part of the GAZYVA infusion-related reaction. Consider withholding antihypertensive treatments for 12 hours prior to, during each GAZYVA infusion, and for the first hour after administration until blood pressure is stable. For patients at increased risk of hypertensive crisis, consider the benefits versus the risks of withholding their antihypertensive medication as is suggested here.

5.4 Hypersensitivity Reactions Including Serum Sickness

Hypersensitivity reactions have been reported in patients treated with GAZYVA. Signs of immediate-onset hypersensitivity included dyspnea, bronchospasm, hypotension, urticaria and tachycardia. Late-onset hypersensitivity diagnosed as serum sickness has also been reported, with symptoms that include chest pain, diffuse arthralgia and fever. Hypersensitivity reactions may be difficult to clinically distinguish from infusion-related reactions. However, hypersensitivity very rarely occurs with the first infusion and, when observed, often occurs after previous exposure.

If a hypersensitivity reaction is suspected during or after an infusion, stop the infusion and permanently discontinue treatment. GAZYVA is contraindicated in patients with known hypersensitivity reactions to GAZYVA, including serum sickness with prior obinutuzumab use [see *Contraindications (4)*].

5.5 Tumor Lysis Syndrome

Tumor lysis syndrome (TLS), including fatal cases, has been reported in patients receiving GAZYVA. Patients with high tumor burden, high circulating lymphocyte count ($> 25 \times 10^9/L$) or renal impairment are at greater risk for TLS.

Administer appropriate tumor lysis prophylaxis with anti-hyperuricemics (e.g., allopurinol or rasburicase) and hydration prior to the infusion of GAZYVA for patients at risk for TLS [see *Dosage and Administration (2.4)*]. During the initial days of GAZYVA treatment, monitor the laboratory parameters of patients considered at risk for TLS. For treatment of TLS, correct electrolyte abnormalities, monitor renal function and fluid balance, and administer supportive care, including dialysis as indicated.

5.6 Infections

Fatal and serious bacterial, fungal, and new or reactivated viral infections can occur during and following GAZYVA therapy. When GAZYVA is administered with chemotherapy followed by GAZYVA monotherapy as in the GALLIUM study, Grade 3 to 5 infections have been reported in up

to 8% of patients during combination therapy, up to 13% of patients during monotherapy, and up to 8% of patients after treatment [see *Adverse Reactions (6.1)*].

In GALLIUM, more Grade 3 to 5 infections were reported in the recipients of GAZYVA and bendamustine (117/410 patients, 29%) as compared to GAZYVA plus CHOP or CVP (43/281 patients, 15%). More fatal infections were reported in patients treated with GAZYVA and bendamustine (3%), as compared to GAZYVA plus CHOP or CVP (< 1%), including during the monotherapy phase and after completion of treatment.

Do not administer GAZYVA to patients with an active infection. Patients with a history of recurring or chronic infections may be at increased risk of infection.

5.7 Neutropenia

Severe and life-threatening neutropenia, including febrile neutropenia, has been reported during treatment with GAZYVA. Monitor patients with Grade 3 to 4 neutropenia frequently with regular laboratory tests until resolution. Anticipate, evaluate, and treat any symptoms or signs of developing infection. Consider dose delays for Grade 3 or 4 neutropenia. Consider administration of granulocyte colony-stimulating factors (GCSF) in patients with Grade 3 or 4 neutropenia.

Neutropenia can also be of late onset (occurring more than 28 days after completion of treatment) and/or prolonged (lasting longer than 28 days). Patients with severe and long lasting (> 1 week) neutropenia are strongly recommended to receive antimicrobial prophylaxis until resolution of neutropenia to Grade 1 or 2. Consider antiviral and antifungal prophylaxis.

5.8 Thrombocytopenia

Severe and life-threatening thrombocytopenia has been reported during treatment with GAZYVA in combination with chemotherapy. Fatal hemorrhagic events have been reported in patients with NHL and CLL treated with GAZYVA in combination with chemotherapy, including during Cycle 1.

Monitor all patients frequently for thrombocytopenia and hemorrhagic events, especially during the first cycle. In patients with Grade 3 or 4 thrombocytopenia, monitor platelet counts more frequently until resolution and consider dose delays of GAZYVA and chemotherapy or dose reductions of chemotherapy. Transfusion of blood products (i.e., platelet transfusion) may be necessary. Consider withholding concomitant medications, which may increase bleeding risk (platelet inhibitors, anticoagulants), especially during the first cycle.

5.9 Immunization

The safety and efficacy of immunization with live or attenuated viral vaccines during or following GAZYVA therapy have not been studied. Immunization with live virus vaccines is not recommended during treatment and until B-cell recovery.

5.10 Embryo-Fetal Toxicity

Based on its mechanism of action and findings in animals, GAZYVA can cause B-cell depletion in infants exposed to obinutuzumab in-utero. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception while receiving GAZYVA and for 6 months after the last dose [see *Use in Specific Populations (8.1, 8.3)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Hepatitis B virus reactivation [see *Warnings and Precautions (5.1)*]
- Progressive multifocal leukoencephalopathy [see *Warnings and Precautions (5.2)*]
- Infusion-related reactions [see *Warnings and Precautions (5.3)*]
- Hypersensitivity reactions including serum sickness [see *Warnings and Precautions (5.4)*]

- Tumor lysis syndrome [see Warnings and Precautions (5.5)]
- Infections [see Warnings and Precautions (5.6)]
- Neutropenia [see Warnings and Precautions (5.7)]
- Thrombocytopenia [see Warnings and Precautions (5.8)]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Chronic Lymphocytic Leukemia

The data below are based on a safety population of 773 previously untreated patients with CLL in the CLL11 study. Patients were treated with chlorambucil alone, GAZYVA in combination with chlorambucil, or rituximab product in combination with chlorambucil. The Stage 1 analysis compared GAZYVA in combination with chlorambucil vs. chlorambucil alone and Stage 2 compared GAZYVA in combination with chlorambucil vs. rituximab product in combination with chlorambucil. Adverse reactions rates and laboratory abnormalities from the Stage 2 phase are presented below and are consistent with the rates in Stage 1. In addition to the adverse reactions observed in Stage 2, in Stage 1, back pain (5% vs. 2%), anemia (12% vs. 10%) and cough (10% vs. 7%) were observed at a higher incidence in the obinutuzumab treated patients. The incidence of Grade 3 to 4 back pain (< 1% vs. 0%), cough (0% vs. < 1%) and anemia (5% vs. 4%) was similar in both treatment arms. With regard to laboratory abnormalities, in Stage 1 hyperkalemia (33% vs. 18%), creatinine increased (30% vs. 20%) and alkaline phosphatase increased (18% vs. 11%) were observed at a higher incidence in patients treated with obinutuzumab with similar incidences of Grade 3 to 4 abnormalities between the two arms.

Patients received three 1,000 mg doses of GAZYVA on the first cycle and a single dose of 1,000 mg once every 28 days for 5 additional cycles in combination with chlorambucil (6 cycles of 28 days each in total). In the last 140 patients enrolled, the first dose of GAZYVA was split between day 1 (100 mg) and day 2 (900 mg) [see Dosage and Administration (2.2)]. In total, 81% of patients received all 6 cycles (of 28 days each) of GAZYVA-based therapy.

Adverse reactions in $\geq 10\%$ of patients in the GAZYVA containing arm were infusion-related reactions, neutropenia, thrombocytopenia, and diarrhea. The most common Grade 3 to 4 adverse reactions (incidence $\geq 10\%$) in the GAZYVA containing arm were neutropenia, infusion-related reactions, and thrombocytopenia.

Table 4 Adverse Reactions (Incidence $\geq 5\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with CLL (Stage 2)

Body System Adverse Reactions	GAZYVA + Chlorambucil n = 336		Rituximab product + Chlorambucil n = 321	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Injury, Poisoning and Procedural Complications				
Infusion-Related Reaction	66	20	38	4
Blood and Lymphatic System Disorders^a				
Neutropenia	38	33	32	28
Thrombocytopenia	14	10	7	3
Gastrointestinal Disorders				
Diarrhea	10	2	8	< 1
Constipation	8	0	5	0
General Disorders and Administration Site Conditions				

Body System Adverse Reactions	GAZYVA + Chlorambucil n = 336		Rituximab product + Chlorambucil n = 321	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Pyrexia	9	< 1	7	< 1
Infections and Infestations				
Nasopharyngitis	6	< 1	3	0
Urinary Tract Infection	5	1	2	< 1

^a Adverse reactions reported under “Blood and lymphatic system disorders” reflect those reported by investigator as clinically significant.

Table 5 Post-Baseline Laboratory Abnormalities (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with CLL (Stage 2)

Laboratory Abnormalities	GAZYVA + Chlorambucil n = 336		Rituximab product + Chlorambucil n = 321	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Hematology				
Leukopenia	84	35	62	16
Lymphopenia	80	39	50	16
Neutropenia	76	46	69	41
Thrombocytopenia	48	13	40	8
Anemia	39	10	37	10
Chemistry				
Hypocalcemia	37	3	32	< 1
ALT increased	28	2	21	1
AST increased	27	2	21	< 1
Hyponatremia	26	7	18	2
Hypoalbuminemia	23	< 1	16	< 1
Hypokalemia	14	1	10	< 1

Non-Hodgkin Lymphoma

GADOLIN

The GADOLIN study evaluated safety in 407 patients with relapsed or refractory NHL, including FL (81%), small lymphocytic lymphoma and marginal zone lymphoma (a disease for which GAZYVA is not indicated), who did not respond to or progressed within 6 months of treatment with rituximab product or a rituximab product-containing regimen. In the population of patients with FL, the profile of adverse reactions was consistent with the overall NHL population. Patients received either GAZYVA in combination with bendamustine (204 patients), followed by GAZYVA monotherapy in patients that had not progressed, or bendamustine alone (203 patients).

Patients randomized to the GAZYVA + bendamustine arm received three weekly 1,000 mg doses of GAZYVA in the first cycle and a single dose of 1,000 mg once every 28 days for 5 additional cycles, in combination with bendamustine 90 mg/m² intravenously on Days 1 and 2 in all 6 cycles. Patients who did not progress on the combination received a single 1,000 mg dose of GAZYVA monotherapy every two months until progression or for a maximum of two years. The control arm received bendamustine 120 mg/m² on Days 1 and 2 of each cycle for 6 cycles, with a cycle length of 28 days. In the GAZYVA arm, 78% of patients received 6 cycles of bendamustine and 82% received their full 6 cycles of GAZYVA; 72 (46%) of the 158 patients who began GAZYVA monotherapy received all planned doses. In the control arm, 72% of patients received 6 cycles of bendamustine.

Serious adverse reactions occurred in 45% of the GAZYVA arm and 37% of the bendamustine-only arm. Fatal adverse reactions within 90 days of treatment occurred in 3.4% and 2.5%, respectively. During treatment and follow-up combined, fatal adverse reactions occurred in 10% of GAZYVA recipients and in 7.4% of recipients of bendamustine alone, with infection and second primary malignancies being the leading causes.

Dose modification due to adverse reactions occurred in 50% of the GAZYVA arm and 42% of the control arm, and discontinuation of any study drug due to adverse reactions occurred in 20% and 17%, respectively.

Table 6 presents selected adverse reactions in GADOLIN. The most common adverse reactions (incidence $\geq 20\%$) in GAZYVA recipients included infusion-related reactions, fatigue, neutropenia, cough, upper respiratory tract infections, and musculoskeletal pain.

Table 6 Adverse Reactions (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with Relapsed or Refractory NHL (GADOLIN)

Body System Adverse Reactions ^{a, b}	GAZYVA + Bendamustine followed by GAZYVA monotherapy n = 204		Bendamustine n = 203	
	All Grades %	Grades 3 to 5 %	All Grades %	Grades 3 to 5 %
Procedural Complications				
Infusion-Related Reaction ^c	67	11	63	5
General Disorders				
Fatigue	40	3	36	3
Pyrexia	19	1	15	1
Blood and Lymphatic System Disorders				
Neutropenia	37	35 ^d	29	27
Infections and Infestations				
Upper Respiratory Tract Infection	36	3	23	1
Respiratory Tract Infection, Unspecified	14	1	8	0
Urinary Tract Infection	13	3	7	0
Respiratory, Thoracic and Mediastinal Disorders				
Cough	31	<1	21	0
Musculoskeletal and Connective Tissue Disorders				
Musculoskeletal Pain	28	1	20	0
Arthralgia	12	<1	5	0
Skin and Subcutaneous Tissue Disorders				
Rash	17	<1	14	<1
Pruritus	11	0	6	0

^a Includes adverse reactions reported throughout study treatment and follow-up.

^b Includes grouped terms.

^c Except where noted, individual events that meet the definition of “infusion-related reaction” are excluded from Table 6 above, as they are included in the grouped term “Infusion-Related Reaction”.

^d Includes 1 fatal event.

Infusion-related reactions are defined as any related adverse reaction that occurred during or within 24 hours of infusion.

Fatigue includes fatigue, lethargy, asthenia.

Pyrexia includes pyrexia, hyperthermia, body temperature increased.

Cough includes cough, productive cough, upper-airway cough syndrome.

Neutropenia includes neutropenia, agranulocytosis, granulocytopenia, neutrophil count decreased.

Upper respiratory tract infection includes upper respiratory tract congestion, upper respiratory tract inflammation, upper respiratory fungal infection, rhinovirus infection, and all terms containing: upper respiratory tract infection, laryngitis, nasopharyngitis, pharyngitis, rhinitis, tonsillitis, and sinusitis with the exception of sinobronchitis.

Respiratory tract infection unspecified includes respiratory tract infection, respiratory tract infection viral, influenza, influenza-like illness, sinobronchitis, respiratory syncytial virus infection.

Urinary tract infection includes all terms containing: urinary tract infection, cystitis, pyelonephritis.

Musculoskeletal pain includes non-cardiac chest pain, bone pain, spinal pain, myalgia, back pain, neck pain, musculoskeletal discomfort, pain in extremity, and all terms containing “musculoskeletal pain”.

Rash includes drug eruption, skin reaction, all terms containing “rash”, urticaria, and selected terms containing “dermatitis”.

Pruritus includes pruritus, pruritus generalized.

Other clinically relevant adverse reactions (incidence $< 10\%$ and $\geq 2\%$ greater in the GAZYVA arm) included:

- *Blood and lymphatic system disorders*: febrile neutropenia (6%)
- *Infection*: sepsis (7%)

During GAZYVA monotherapy (158 patients), adverse reactions in $\geq 10\%$ of patients included upper and lower respiratory tract infections, cough, neutropenia, musculoskeletal pain, fatigue, diarrhea, rash, and urinary tract infection.

Table 7 presents selected new or worsening laboratory abnormalities in the GADOLIN trial.

Table 7 New or Worsening Laboratory Abnormalities (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm^a) in Patients with Relapsed or Refractory NHL (GADOLIN)

Laboratory Abnormalities	GAZYVA + Bendamustine followed by GAZYVA monotherapy n = 204		Bendamustine n = 203	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Hematology				
Lymphopenia	97	92	96	84
Leukopenia	84	47	87	34
Neutropenia	76	53	75	42
Chemistry				
Hypophosphatemia	41	8	38	7
Hypocalcemia	39	3	24	1
ALT/SGPT increased	36	2	31	3
Alkaline phosphatase increased	27	0	23	0
Hyperbilirubinemia	21	2	17	2
Hyperkalemia	20	3	18	0

^a Two percent difference in either any-grade or Grade 3 to 4 laboratory abnormalities.

In the GAZYVA monotherapy phase, new or worsening grade 3 or 4 abnormalities included neutropenia in 25% of patients (Grade 4, 10%) and lymphopenia in 23% (Grade 4, 5%).

GALLIUM

A randomized, open-label multicenter trial (GALLIUM) evaluated the safety of GAZYVA as compared to rituximab product in 1385 patients with previously untreated follicular lymphoma (86%) or marginal zone lymphoma (14%). Patients received chemotherapy (bendamustine, CHOP, or CVP) combined with either GAZYVA (691 patients) or rituximab product (694 patients), followed in responding patients by GAZYVA or rituximab product monotherapy every two months until disease progression or for a maximum of two years. The study excluded patients having an absolute neutrophil count (ANC) $< 1500 / \mu\text{L}$, platelets $< 75,000 / \mu\text{L}$, CLcr $< 40 \text{ mL/min}$ and, unless attributable to lymphoma, hepatic transaminases $> 2.5 \times$ upper limit of normal.

The median age was 60 (range: 23-88), 47% were male, 82% were white, and 97% had an ECOG performance status of 0 or 1. The chemotherapy was bendamustine in 59%, CHOP in 31% and CVP in 10% of patients. Following combination therapy, 624 patients (90%) in the GAZYVA arm and 612 patients (88%) in the rituximab product arm received monotherapy.

Serious adverse reactions occurred in 50% of patients on the GAZYVA arm and 43% of patients on the rituximab product arm. Fatal adverse reactions were reported during treatment in 3% in the GAZYVA arm and 2% in the rituximab product arm, most often from infections in the GAZYVA arm. During treatment and follow-up combined, fatal adverse reactions were reported in 5% of the GAZYVA arm and 4% of the rituximab product arm, with infections and second malignancies being leading causes. In the GAZYVA arm, fatal infections occurred in 2% of patients compared to $< 1\%$ in the rituximab product arm.

During combination therapy, 93% of patients received all treatment cycles in the GAZYVA arm, and 92% received all treatment cycles in the rituximab product arm. Of the responding patients who began

monotherapy with GAZYVA or rituximab product, 76% and 73%, respectively, completed the full course. Dose modification due to adverse reactions occurred in 74% of the GAZYVA arm and 63% of the rituximab product arm throughout study treatment, and discontinuation of any study drug due to adverse reactions occurred in 18% and 15%, respectively.

Throughout treatment and follow-up, the most common adverse reactions (incidence $\geq 20\%$) observed at least 2% more in the GAZYVA arm included infusion-related reactions, neutropenia, upper respiratory tract infections, constipation and diarrhea (Table 8). Neutropenia, infusion-related reactions, febrile neutropenia and thrombocytopenia were the most common Grade 3 to 5 adverse reactions (incidence $\geq 5\%$) observed more frequently in the GAZYVA arm.

Table 8 Adverse Reactions (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with Previously Untreated NHL (GALLIUM)

Body System Adverse Reactions ^{a, b}	GAZYVA + chemotherapy followed by GAZYVA monotherapy n = 691		Rituximab product + chemotherapy followed by rituximab product monotherapy n = 694	
	All Grades %	Grades 3 to 5 %	All Grades %	Grades 3 to 5 %
Injury, Poisoning and Procedural Complications				
Infusion-Related Reaction ^c	72	12	60	8
Blood and Lymphatic System Disorders				
Neutropenia ^d	53	49	47	41
Thrombocytopenia ^d	14	7	8	3
Infections and Infestations				
Upper Respiratory Tract Infection	50	3	43	1
Herpesvirus Infection	18	3	14	1
Pneumonia	14	7	12	6
Respiratory, Thoracic and Mediastinal Disorders				
Cough	35	< 1	28	< 1
Gastrointestinal Disorders				
Constipation	32	< 1	29	< 1
Diarrhea	30	3	26	2
Nervous System Disorders				
Headache	18	< 1	15	< 1
Musculoskeletal and Connective Tissue Disorders				
Arthralgia	16	0	14	< 1
Psychiatric Disorders				
Insomnia	15	< 1	12	< 1
Metabolism and Nutrition Disorders				
Decreased Appetite	14	< 1	12	< 1
Skin and Subcutaneous Tissue Disorders				
Pruritus	11	< 1	9	0

^a Includes adverse reactions reported throughout study treatment and follow-up.

^b Includes grouped terms.

^c Except where noted, individual events that meet the definition of “infusion-related reaction” are excluded from Table 8 above, as they are already included in the grouped term “Infusion-Related Reaction”. The most common individual terms within the grouped term “Infusion-Related Reaction” in decreasing order of frequency are nausea, chills, pyrexia and vomiting.

^d Includes adverse reactions reported as infusion-related reactions.

Infusion-related reactions are defined as any related adverse reaction that occurred during or within 24 hours of infusion.

Neutropenia includes neutropenia, agranulocytosis, granulocytopenia, and neutrophil count decreased.

Febrile neutropenia includes febrile neutropenia, neutropenic infection, neutropenic sepsis, and febrile bone marrow aplasia.

Thrombocytopenia includes thrombocytopenia and platelet count decreased.

Upper respiratory tract infection includes upper respiratory tract congestion, upper respiratory tract inflammation, upper respiratory tract infection, rhinovirus infection, and all terms containing: laryngitis, nasopharyngitis, pharyngitis, rhinitis, tonsillitis, and sinusitis with the exception of sinobronchitis.

Herpesvirus infection includes all terms containing “herpes” or “varicella.”

Pneumonia includes all terms containing “pneumonia,” bacterial, pneumonia haemophilus, pneumonia pneumococcal, pneumonia fungal, pneumocystis jirovecii infection, lung infection, and lung infiltration.

Diarrhea includes diarrhea, defecation urgency, frequent bowel movement, and all terms containing “gastroenteritis”.

Headache includes all terms containing “headache” and migraine.

Insomnia includes all terms containing “insomnia” and sleep disorder.

Pruritus includes pruritus and pruritus generalized.

During the monotherapy period, the common adverse reactions (incidence $\geq 10\%$) observed at least 2% more with GAZYVA were upper respiratory tract infection (40%), cough (23%), musculoskeletal pain (20%), neutropenia (19%), and herpesvirus infection (13%).

Table 9 summarizes treatment-emergent laboratory abnormalities during treatment and follow-up. The Grade 3 to 4 abnormalities reported at least 2% more in the GAZYVA arm were lymphopenia, leukopenia, neutropenia, thrombocytopenia, and hyperuricemia. Patients in the GAZYVA arm, as compared to the rituximab product arm, had higher incidences of Grade 4 neutropenia (38% vs. 30%, respectively), Grade 4 lymphopenia (33% vs. 22%), and Grade 4 leukopenia (17% vs. 12%).

Table 9 New or Worsening Laboratory Abnormalities (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with Previously Untreated NHL (GALLIUM)

Laboratory Abnormalities ^a	GAZYVA+ chemotherapy followed by GAZYVA monotherapy n = 691		Rituximab product + chemotherapy followed by rituximab product monotherapy n = 694	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Hematology				
Lymphopenia	97	83	95	67
Leukopenia	92	49	89	39
Neutropenia	84	59	76	50
Thrombocytopenia	68	11	50	4
Chemistry				
ALT increased	50	3	43	2
AST increased	44	1	41	1
Hypophosphatemia	36	5	33	5
Hypoalbuminemia	33	1	25	1
Hypoproteinemia	32	0	30	0
Hypocalcemia	32	1	26	1
Hyperuricemia	28	28	22	22
Hyponatremia	26	4	20	3
Hyperkalemia	23	1	17	1
Hypernatremia	16	< 1	13	0

^a Includes lab abnormalities, reported throughout treatment and follow-up, that were new or worsening, or worsening from baseline unknown.

In the monotherapy phase, new-onset Grade 3 or 4 neutropenia was reported in 21% of patients in the GAZYVA arm (Grade 4, 10%) and 17% of patients in the rituximab product arm (Grade 4, 9%).

Infusion-Related Reactions

Chronic Lymphocytic Leukemia

The incidence of infusion-related reactions (IRRs) in the CLL11 study was 65% with the first infusion of GAZYVA. The incidence of Grade 3 or 4 IRRs was 20% with 7% of patients discontinuing therapy. The incidence of reactions with subsequent infusions was 3% with the second 1,000 mg and < 1% thereafter. No Grade 3 or 4 IRRs were reported beyond the first 1,000 mg infused.

Of the first 53 patients receiving GAZYVA in CLL11, 47 (89%) experienced an IRR. After this experience, study protocol modifications were made to require pre-medication with a corticosteroid, antihistamine, and acetaminophen. The first dose was also divided into two infusions (100 mg on day 1 and 900 mg on day 2). For the 140 patients for whom these mitigation measures were implemented, 74 patients (53%) experienced a reaction with the first 1,000 mg (64 patients on day 1, 3 patients on day 2, and 7 patients on both days) and < 3% thereafter [see *Dosage and Administration (2.2)*].

Non-Hodgkin Lymphoma

Overall, 67% of patients in the GADOLIN study experienced an IRR (all grades) during treatment with GAZYVA in combination with bendamustine. The incidence of Grade 3 to 4 IRRs in GADOLIN was 11%. In Cycle 1, the incidence of IRRs (all grades) was 53% in patients receiving GAZYVA in combination with bendamustine of which 34 (9%) were Grade 3 to 4 in severity. In patients receiving GAZYVA in combination with bendamustine, the incidence of IRRs was highest on Day 1 (37%), and gradually decreased on Days 2, 8 and 15 (23%, 6% and 4%, respectively).

During Cycle 2, the incidence of IRRs was 24% in patients receiving GAZYVA in combination with bendamustine and decreased with subsequent cycles.

During GAZYVA monotherapy in GADOLIN, IRRs (all grades) were observed in 8% of patients. One Grade 3 and no Grade 4 IRRs were reported during GAZYVA monotherapy.

Overall, 2% of patients in GADOLIN experienced an IRR leading to discontinuation of GAZYVA.

In GALLIUM, 72% of patients in the GAZYVA treated arm experienced an IRR (all grades). The incidence of Grade 3 to 4 IRRs for these patients was 12%. In Cycle 1, the incidence of IRRs (all grades) was 62% in the GAZYVA treated arm with Grade 3 to 4 IRRs reported in 10%. The incidence of IRRs (all grades) was highest on Day 1 (60%) and decreased on Days 8 and 15 (9% and 6%, respectively).

During Cycle 2, the incidence of IRRs (all grades) in the GAZYVA treated arm was 13% and decreased with subsequent cycles.

During GAZYVA monotherapy treatment in GALLIUM, IRRs (all grades) were observed in 9% of patients.

Overall, 1% of patients in GALLIUM experienced an IRR leading to discontinuation of GAZYVA.

Neutropenia

Chronic Lymphocytic Leukemia

The incidence of neutropenia reported as an adverse reaction in CLL11 was 38% in the GAZYVA treated arm and 32% in the rituximab product treated arm, with the incidence of serious adverse reactions being 1% and < 1%, respectively (Table 4). Cases of late-onset neutropenia (occurring 28 days after completion of treatment or later) were 16% in the GAZYVA treated arm and 12% in the rituximab product treated arm.

Non-Hodgkin Lymphoma

The incidence of neutropenia in GADOLIN was higher in the GAZYVA plus bendamustine arm (37%) compared to the arm treated with bendamustine alone (30%). Cases of prolonged neutropenia (3%) and late onset neutropenia (8%) were also reported in the GAZYVA plus bendamustine arm. The incidence of neutropenia was higher during treatment with GAZYVA in combination with bendamustine (30%) compared to the GAZYVA monotherapy treatment phase (13%).

The incidence of neutropenia in GALLIUM was higher in the GAZYVA treated arm (53%) compared to the rituximab product treated arm (47%). Cases of prolonged neutropenia (1%) and late onset neutropenia (4%) were also reported in the GAZYVA treated arm. The incidence of neutropenia was

higher during treatment with GAZYVA in combination with chemotherapy (45%) compared to the GAZYVA monotherapy treatment phase (20%).

Infection

Chronic Lymphocytic Leukemia

The incidence of infections was similar between GAZYVA and rituximab product treated arms. Thirty-eight percent of patients in the GAZYVA treated arm and 37% in the rituximab product treated arm experienced an infection, with Grade 3 to 4 rates being 11% and 13%, respectively. Fatal events were reported in 1% of patients in both arms.

Non-Hodgkin Lymphoma

The incidence of infection in GADOLIN was 68% in the GAZYVA plus bendamustine arm and 59% in the bendamustine arm, with Grade 3 to 4 events reported in 20% and 16%, respectively. Fatal events were reported in 3% of patients in the GAZYVA plus bendamustine arm and 3% in the bendamustine arm.

The incidence of infections in GALLIUM was 82% in the GAZYVA treated arm and 73% in the rituximab product treated arm, with Grade 3 to 4 events reported in 21% and 17%, respectively. In the GAZYVA arm, fatal infections occurred in 2% of patients compared to <1% in the rituximab product arm.

The incidence of Grade 3 to 4 infections in the GAZYVA and rituximab product treated arms was lower in patients receiving GCSF prophylaxis (14%; 16%) compared with patients not receiving GCSF prophylaxis (24%; 18%). The incidence of fatal infections in patients receiving GCSF prophylaxis in the GAZYVA and rituximab product treated arms was 2% and 0%, respectively, and was 2% and < 1% in patients not receiving GCSF prophylaxis.

Thrombocytopenia

Chronic Lymphocytic Leukemia

The overall incidence of thrombocytopenia reported as an adverse reaction was higher in the GAZYVA treated arm (14%) compared to the rituximab product treated arm (7%), with the incidence of Grade 3 to 4 events being 10% and 3%, respectively (Table 4). The difference in incidences between the treatment arms is driven by events occurring during the first cycle. The incidence of thrombocytopenia (all grades) in the first cycle was 11% in the GAZYVA and 3% in the rituximab product treated arms, with Grade 3 to 4 rates being 8% and 2%, respectively. Four percent of patients in the GAZYVA treated arm experienced acute thrombocytopenia (occurring within 24 hours after the GAZYVA infusion).

The overall incidence of hemorrhagic events and the number of fatal hemorrhagic events were similar between the treatment arms, with 3 in the rituximab product and 4 in the GAZYVA treated arms. However, all fatal hemorrhagic events in patients treated with GAZYVA occurred in Cycle 1.

Non-Hodgkin Lymphoma

The incidence of thrombocytopenia in GADOLIN was lower in the GAZYVA plus bendamustine arm (15%) compared to the arm treated with bendamustine alone (25%). The incidence of hemorrhagic events in GAZYVA plus bendamustine treated patients compared to bendamustine alone was 12% and 11%, respectively. Grade 3 to 4 hemorrhagic events were similar in both treatment arms (4% in the GAZYVA plus bendamustine arm and 2% in the bendamustine arm).

In GALLIUM, thrombocytopenia was reported as an adverse reaction in 14% of the GAZYVA treated arm and 8% of the rituximab product treated arm, with the incidence of Grade 3 to 4 events being 7% and 3%, respectively. The difference in incidences between the treatment arms is driven by events occurring during the first cycle. The incidence of thrombocytopenia (all grades) in the first cycle was

9% in the GAZYVA and 3% in the rituximab product treated arms, with Grade 3 to 4 rates being 5% and 1%, respectively. In GALLIUM, both treatment arms had a 12% overall incidence of hemorrhagic events and a < 1% incidence of fatal hemorrhagic events.

Tumor Lysis Syndrome

The incidence of Grade 3 or 4 tumor lysis syndrome in GAZYVA treated patients was 2% in CLL11, 0.5% in GADOLIN and 0.9% in GALLIUM.

Musculoskeletal Disorders

Chronic Lymphocytic Leukemia

Adverse reactions related to musculoskeletal disorders (all events from the body system), including pain, have been reported in the GAZYVA treated arm with higher incidence than in the rituximab product treated arm (18% vs. 15%).

Non-Hodgkin Lymphoma

In GADOLIN, adverse reactions related to musculoskeletal disorders (all events from the body system), including pain, have been reported in the GAZYVA plus bendamustine treated arm with higher incidence than in the bendamustine alone arm (44% vs. 30%).

In GALLIUM, musculoskeletal disorders were reported in 54% of patients in the GAZYVA treated arm and 49% of patients in the rituximab product treated arm.

Liver Enzyme Elevations

Hepatic enzyme elevations have occurred in CLL patients who received GAZYVA in clinical trials and had normal baseline hepatic enzyme levels (AST, ALT and ALP). The events occurred most frequently within 24–48 hours of the first infusion. In some patients, elevations in liver enzymes were observed concurrently with infusion-related reactions or tumor lysis syndrome. In the CLL11 study, there was no clinically meaningful difference in overall hepatotoxicity adverse reactions between all arms (4% of patients in the GAZYVA treated arm). Medications commonly used to prevent infusion-related reactions (e.g., acetaminophen) may also be implicated in these events. Monitor liver function tests during treatment, especially during the first cycle. Consider treatment interruption or discontinuation for hepatotoxicity.

Gastrointestinal Perforation

Cases of gastrointestinal perforation have been reported in patients receiving GAZYVA, mainly in NHL.

Worsening of Pre-existing Cardiac Conditions

Fatal cardiac events have been reported in patients treated with GAZYVA.

6.2 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies in the studies described below with the incidence of antibodies in other studies or to other products may be misleading.

Seven percent (18/271) of patients with CLL tested positive for anti-GAZYVA antibodies at one or more time points in CLL11. No patients developed anti-GAZYVA antibodies during or following GAZYVA treatment in GADOLIN, while 1 patient (1/564, 0.2%) developed anti-GAZYVA antibodies in GALLIUM. Neutralizing activity of anti-GAZYVA antibodies has not been assessed.

6.3 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of GAZYVA. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- *Immune/Autoimmune Events*: Serum sickness

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings from animal studies and its mechanism of action, GAZYVA can cause fetal B-cell depletion [see *Clinical Pharmacology (12.1)*]. There are no data with GAZYVA use in pregnant women to inform a drug-associated risk. Monoclonal antibodies are transferred across the placenta. In animal reproduction studies, weekly intravenous administration of obinutuzumab to pregnant cynomolgus monkeys from day 20 of pregnancy until parturition which includes the period of organogenesis at doses with exposures up to 2.4 times the exposure at the clinical dose of 1,000 mg monthly produced opportunistic infections and immune complex mediated hypersensitivity reactions. No embryo-toxic or teratogenic effects were observed in the monkeys (see *Data*). Advise pregnant women of the potential risk to the fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown; however, the estimated background risk in the U.S. general population of major birth defects is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregnancies.

Clinical Considerations

Fetal/Neonatal Adverse Reactions

GAZYVA is likely to cause fetal B-cell depletion (see *Data*). Avoid administering live vaccines to neonates and infants exposed to GAZYVA in utero until B-cell recovery occurs [see *Warnings and Precautions (5.8)* and *Clinical Pharmacology (12.2)*].

Data

Animal Data

In a pre- and post-natal development study, pregnant cynomolgus monkeys received weekly intravenous doses of 25 or 50 mg/kg obinutuzumab from day 20 of pregnancy until parturition, which includes the period of organogenesis. The high dose results in an exposure (AUC) that is 2.4 times the exposure in patients with CLL at the recommended label dose. There were no embryo-toxic or teratogenic effects in animals. Secondary opportunistic infections, immune complex mediated hypersensitivity reactions, or a combination of both were observed in exposed dams. When first measured on day 28 postpartum, obinutuzumab was detected in offspring at levels in the range of maternal serum levels on the same day, and B-cells were completely depleted. The B-cell counts returned to normal levels, and immunologic function was restored within 6 months after birth.

Obinutuzumab was measured in the milk of lactating cynomolgus monkeys on day 28 postpartum after weekly intravenous administration from day 20 of pregnancy until parturition. Concentrations in milk were approximately 0.04% and 0.13% of concentrations in maternal serum in the 25 and 50 mg/kg groups, respectively.

8.2 Lactation

Risk Summary

There is no information regarding the presence of GAZYVA in human milk, the effects on the breastfed child, or the effects on milk production. However, low levels of obinutuzumab were present

in the milk of lactating cynomolgus monkeys [see *Use in Specific Populations (8.1)*]. Human IgG is known to be present in human milk. Because of the potential of serious adverse reactions in the breastfed child, advise women not to breastfeed during treatment with GAZYVA and for 6 months after the last dose.

8.3 Females and Males of Reproductive Potential

Contraception

GAZYVA can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

Females

Advise females of reproductive potential to use effective contraception during treatment with GAZYVA and for 6 months after the last dose.

8.4 Pediatric Use

The safety and effectiveness of GAZYVA in pediatric patients have not been established.

8.5 Geriatric Use

Chronic Lymphocytic Leukemia

Of 336 patients with previously untreated CLL who received GAZYVA in combination with chlorambucil, 81% were 65 years and older, while 46% were 75 and older. Of the patients 75 years and older, 46% experienced serious adverse reactions and 7% experienced adverse reactions leading to death. Of the patients younger than 75, 33% experienced a serious adverse reaction and 2% an adverse reaction leading to death. No significant differences in efficacy were observed between younger and older patients [see *Clinical Studies (14.1)*].

Non-Hodgkin Lymphoma

Of 204 patients in GADOLIN with relapsed or refractory NHL treated with GAZYVA plus bendamustine, 44% were 65 and over, while 14% were 75 and over. In patients 65 and over, 55% of patients experienced serious adverse reactions and 28% experienced adverse reactions leading to treatment withdrawal while in patients under 65, 37% and 14% experienced serious adverse reactions and adverse reactions leading to treatment withdrawal, respectively. No clinically meaningful differences in efficacy were observed between these patients and younger patients in GADOLIN.

Of the 691 patients in GALLIUM treated with GAZYVA plus chemotherapy as first-line therapy, 33% were 65 and over, while 7% were 75 and over. Of patients 65 and over, 63% experienced serious adverse reactions and 26% experienced adverse reactions leading to treatment withdrawal, while in patients under 65, 43% experienced serious adverse reactions and 13% had an adverse reaction leading to treatment withdrawal. No clinically meaningful differences in efficacy were observed between these patients and younger patients in GALLIUM.

10 OVERDOSAGE

There has been no experience with overdose in human clinical trials. For patients who experience overdose, treatment should consist of immediate interruption or reduction of GAZYVA and supportive therapy.

11 DESCRIPTION

Obinutuzumab is a humanized anti-CD20 monoclonal antibody of the IgG1 subclass. It recognizes a specific epitope of the CD20 molecule found on B cells. The molecular mass of the antibody is approximately 150 kDa.

GAZYVA (obinutuzumab) injection is produced by mammalian cell (CHO) suspension culture. GAZYVA was engineered for reduced fucose content as compared to a typical IgG1 produced in CHO

cells. GAZYVA is a sterile, clear, colorless to slightly brown, preservative-free liquid concentrate for intravenous use. GAZYVA is supplied at a concentration of 25 mg/mL in 1,000 mg single-dose vials. The product is formulated in 20 mM L-histidine/L-histidine hydrochloride, 240 mM trehalose, 0.02% poloxamer 188. The pH is 6.0.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Obinutuzumab is a monoclonal antibody that targets the CD20 antigen expressed on the surface of pre-B and mature B lymphocytes. Upon binding to CD20, obinutuzumab mediates B-cell lysis through (1) engagement of immune effector cells, (2) by directly activating intracellular death signaling pathways (direct cell death), and/or (3) activation of the complement cascade. The immune effector cell mechanisms include antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis.

As an antibody with reduced fucose content, obinutuzumab induces greater ADCC activity than rituximab in vitro using human cancer cell lines. Obinutuzumab also demonstrated an increased ability to induce direct cell death when compared to rituximab. Obinutuzumab binds to FcγRIII using purified proteins with a higher affinity than rituximab. Obinutuzumab and rituximab bind with similar affinity to overlapping epitopes on CD20.

12.2 Pharmacodynamics

In patients with CLL, GAZYVA caused CD19 B-cell depletion (defined as CD19 B cell counts < 0.07 x 10⁹/L). Initial CD19 B cell recovery was observed in some patients approximately 9 months after the last GAZYVA dose. At 18 months of follow-up, some patients remain B cell depleted.

Although the depletion of B cells in the peripheral blood is a measurable pharmacodynamic effect, it is not directly correlated with the depletion of B-cells in solid organs or in malignant deposits. B cell depletion has not been shown to be directly correlated to clinical response.

Cardiac Electrophysiology

The potential effects of GAZYVA on the QTc interval have not been studied.

12.3 Pharmacokinetics

The pharmacokinetic parameters of obinutuzumab after 100 mg on day 1 and 900 mg on day 2 of Cycle 1, 1,000 mg on day 8 and 15 of Cycle 1, and 1,000 mg on day 1 of Cycles 2–6 for CLL and after 1,000 mg on day 1, 8 and 15 of Cycle 1, 1,000 mg on day 1 of Cycles 2-6 or Cycles 2-8, and then 1,000 mg every 2 months for up to 2 years for NHL are provided in Table 10. The dosing regimen is within the linear pharmacokinetic behavior of obinutuzumab.

Table 10 Obinutuzumab Measures of Exposure

PK Measure	CLL ^a	Relapsed or refractory FL ^a	First line FL in combination with chemotherapy	
			GAZYVA + Bendamustine ^a	GAZYVA + CHOP or CVP ^b
C _{max} , µg/mL	466.3 (35)	553.5 (32)	513.4 (28)	676.4 (30)
C _{trough} , µg/mL	192.5 (78)	295 (56)	255 (46)	395 (44)
AUC, µg/mL*day	8701 (51)	11362 (41)	10088 (35)	10723 (37)

Results are presented as geometric mean (% Coefficient of Variation).

^a Induction Cycle 6 of a 28-day cycle;

^b Induction Cycle 8 of a 21-day cycle.

Distribution and Elimination

The elimination of obinutuzumab is comprised of a linear clearance pathway and a time-dependent non-linear clearance pathway. As GAZYVA treatment progresses, the impact of the time-dependent pathway diminishes in a manner suggesting target-mediated drug disposition (TMDD) and saturation of the TMDD at the end of the treatment cycle at the proposed clinical dose regimen. The pharmacokinetic properties of obinutuzumab in patients with CLL and NHL are provided in Table 11.

Table 11 Pharmacokinetic Parameters of Obinutuzumab

	CLL	NHL
Distribution		
Volume of Distribution ^a , L	4.1 (20)	4.3 (21)
Elimination		
Terminal Half-life, days	25.5 (48)	35.3 (35)
Clearance, L/day	0.11 (53)	0.08 (41)

Parameters are presented as geometric mean (% Coefficient of Variation).

^a At steady state.

Specific Populations

Age (median [range]: 63 [22, 89] years) and baseline creatinine clearance (CLCr) (median [range] 84 [22, > 120] mL/min) did not affect the pharmacokinetics of GAZYVA. In patients with CLCr ≤ 30 mL/min, the pharmacokinetics of GAZYVA was unaffected. GAZYVA has not been studied in patients with hepatic impairment.

The volume of distribution and steady-state clearance increased with body weight; however, the expected change in exposure does not warrant a dose modification.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No carcinogenicity or genotoxicity studies have been conducted with obinutuzumab.

No specific studies have been conducted to evaluate potential effects on fertility; however, no adverse effects on male or female reproductive organs were observed in the 26-week repeat-dose toxicity study in cynomolgus monkeys.

14 CLINICAL STUDIES

14.1 Chronic Lymphocytic Leukemia

The efficacy of GAZYVA was evaluated in a three-arm, open-label, active-controlled, randomized, multicenter trial (CLL11; NCT01010061) in 781 patients with previously untreated CD20+ CLL requiring treatment who had coexisting medical conditions or reduced renal function as measured by creatinine clearance (CLCr) < 70 mL/min. Patients with CLCr < 30 mL/min, active infections, positive hepatitis B (HBsAg or anti-HBc positive; patients positive for anti-HBc could be included if hepatitis B viral DNA was not detectable) and hepatitis C serology, or immunization with live virus vaccine within 28 days prior to randomization were excluded from the trial. Patients were treated with chlorambucil control (Arm 1), GAZYVA in combination with chlorambucil (Arm 2), or rituximab product in combination with chlorambucil (Arm 3). The safety and efficacy of GAZYVA was evaluated in a Stage 1 comparison of Arm 1 vs. Arm 2 in 356 patients and a Stage 2 comparison of Arm 2 vs. Arm 3 in 663 patients.

The majority of patients received 1,000 mg of GAZYVA on days 1, 8 and 15 of the first cycle, followed by treatment on the first day of 5 subsequent cycles (total of 6 cycles, 28 days each). The first dose of GAZYVA was divided between day 1 (100 mg) and day 2 (900 mg) [see *Dosage and Administration (2.2)*], which was implemented in 140 patients. Chlorambucil was given orally at 0.5 mg/kg on day 1 and day 15 of all treatment cycles (1 to 6).

In CLL11, the median age was 73 years, 62% were male, and 95% were White. Sixty-five percent had a CLCr < 70 mL/min and 76% had multiple coexisting medical conditions. Twenty-two percent of patients were Binet stage A, 42% were stage B, and 36% were stage C. The median estimated CLCr was 62 mL/min. Eighty-one percent of patients treated with GAZYVA in combination with chlorambucil received all 6 cycles compared to 89% of patients in the rituximab product treated arm and 67% in the chlorambucil alone arm.

In the Stage 1 analysis of CLL11, the median progression-free survival (PFS) in the GAZYVA in combination with chlorambucil arm was 27.2 months and 11.2 months in the chlorambucil alone arm (median observation time 22.8 months) as assessed by independent review and is consistent with investigator-assessed PFS. The median overall survival (OS) was not yet reached with a total of 46 deaths: 22 (9%) in the GAZYVA in combination with chlorambucil arm and 24 (20%) in the chlorambucil arm. The hazard ratio for OS was 0.41 (95% CI: 0.23-0.74).

In the Stage 2 analysis of CLL11, the median PFS was 26.7 months in the GAZYVA arm and 14.9 months in the rituximab product arm with a median observation time of 18.7 months (HR: 0.42, 95% CI: 0.33-0.54, p-value < 0.0001). These results were assessed by independent review and are consistent with investigator-assessed PFS. Minimal residual disease (MRD) was evaluated using allele-specific oligonucleotide polymerase chain reaction (ASO-PCR). The cutoff for a negative status was one CLL cell per 10^4 leukocytes in the sample (i.e., an MRD value of $< 10^{-4}$ was considered negative). Among patients who achieved complete response (CR) and complete response with incomplete marrow recovery (CRi; 94 patients in the GAZYVA arm and 34 patients in the rituximab product arm), 18 patients (19%) had negative MRD in the bone marrow in the GAZYVA arm compared to 2 patients (6%) in the rituximab product arm. Out of the patients who achieved CR and CRi, 39 patients (41%) in the GAZYVA arm, and 4 patients (12%) in the rituximab product arm were MRD negative in peripheral blood samples collected at least 3 months after the end of treatment.

Efficacy results are shown in Table 12 and Figures 1 and 2.

Table 12 Efficacy Results from CLL11

Endpoint	Stage 1 of CLL11		Stage 2 of CLL11	
	GAZYVA + Chlorambucil* n = 238	Chlorambucil n = 118	GAZYVA + Chlorambucil* n = 333	Rituximab product + Chlorambucil n = 330
Median Progression-Free Survival ^a	27.2 months (HR 0.19 [0.14; 0.27], p-value < 0.0001 stratified log-rank test)	11.2 months	26.7 months (HR 0.42 [0.33; 0.54], p-value < 0.0001 stratified log-rank test)	14.9 months
Overall Response Rate ^b	78.2%	33.1%	79.6%	66.3%
Complete Response	28.2%	0	26.1%	8.8%
Complete Response with Incomplete Marrow Recovery	2.5%	1.7%	2.1%	1.5%
Partial Response	45.0%	30.5%	48.6%	54.1%
Nodular Partial Response	2.5%	0.8%	2.7%	1.8%
Median Duration of Response	22.4 months	4.7 months	19.6 months	9.7 months
Overall Survival	HR 0.41 [0.23; 0.74]		Not Yet Mature	

^a As defined by independent review. Investigator-assessed PFS was consistent with data from independent review.

^b Defined as best overall response rate (ORR = CR + CRi + PR + nPR).

*All Stage 1 GClb patients (n = 238) were included in the Stage 2 GClb population (n = 333).

Figure 1
Kaplan-Meier Curve of Overall Survival in Patients with CLL in CLL11 (Stage 1)





Figure 2
Kaplan-Meier Curve of Progression-Free Survival in Patients with CLL in CLL11 (Stage 2)



14.2 Follicular Lymphoma

GADOLIN

The efficacy of GAZYVA was evaluated in GADOLIN (NCT01059630), an open-label, multicenter, randomized study that included 335 patients with follicular lymphoma (FL) who had no response to or have progressed during or within 6 months of rituximab product or a rituximab product-containing regimen. These patients were randomized to receive either bendamustine alone (n = 171) or GAZYVA in combination with bendamustine (n = 164) for 6 cycles, each of 28 days duration. Patients in the GAZYVA plus bendamustine arm who did not have disease progression [patients with a complete response (CR), partial response (PR) or stable disease (SD)] at the end of the 6 cycles continued receiving GAZYVA monotherapy for 2 years. Patients were stratified according to the type of refractoriness to rituximab product (refractory to rituximab product monotherapy versus rituximab product in combination with chemotherapy), the number of prior therapies (≤ 2 versus > 2), and geographic region.

GAZYVA was given by intravenous infusion as a flat dose of 1,000 mg on Days 1, 8 and 15 of Cycle 1, on Day 1 of Cycles 2–6, and then every 2 months until disease progression for up to 2 years. Bendamustine was given intravenously on Days 1 and 2 for all treatment cycles (1–6) at 90 mg/m²/day when given in combination with GAZYVA or 120 mg/m²/day when given alone.

The primary analysis included 321 FL patients, including 166 patients randomized to bendamustine alone and 155 patients randomized to GAZYVA in combination with bendamustine. In the primary analysis, patients had a median age of 63 years, 88% were White and 56% were male. Thirty-four percent had bulky disease (> 6 cm), 15% had at least one B-symptom at baseline and 95% had an ECOG performance status of 0–1 at baseline. The median time since initial diagnosis was 3 years and the median number of prior therapies was 2 (range 1 to 10). Forty-six percent of patients received 1 prior therapy and 33% of patients received 2 prior therapies. Twenty percent of patients were refractory to prior rituximab product monotherapy, 37% of patients were refractory to prior rituximab product plus chemotherapy induction treatment, and 41% of patients were refractory to rituximab product maintenance treatment received following rituximab product plus chemotherapy induction. Seventy-nine percent of patients were refractory to both rituximab product and an alkylating agent during any prior regimen (double refractory).

The major efficacy outcome measure was PFS as determined by an independent review committee (IRC). At the time of the primary analysis, median observation time was 21.1 months. The median PFS in the bendamustine arm was 13.8 months. Median PFS was not reached in the GAZYVA plus bendamustine arm (PFS HR = 0.48, 95% CI: 0.34-0.68; stratified log-rank test p-value < 0.0001). The investigator assessed PFS result was consistent with the IRC-assessed PFS. The median investigator-assessed PFS in the bendamustine arm was 13.7 months and the median in the GAZYVA containing arm was 29.2 months (PFS HR = 0.48, 95% CI: 0.35-0.67; stratified log-rank test p-value < 0.0001).

Efficacy results are summarized in Table 13. The Kaplan-Meier curve for IRC-PFS is shown in Figure 3.

Table 13 Primary Analysis Efficacy Results from GADOLIN^{a, b}

Endpoint	GADOLIN	
	GAZYVA + Bendamustine followed by GAZYVA monotherapy n = 155	Bendamustine n = 166
Median Progression-Free Survival (months)	Not Reached (HR = 0.48 [0.34; 0.68], p-value < 0.0001 by stratified log-rank test)	13.8
Best Overall Response ^c	78.7%	74.7%
Complete Response	15.5%	18.7%
Partial Response	63.2%	56.0%
Median duration of response (months)	Not Reached	11.6


^a Based on FL population.

^b As defined by independent review.

^c Best response of PR or CR within 12 months of study start.


Figure 3

Kaplan-Meier Curve of IRC-Assessed Progression-Free Survival in Patients with FL



The final analysis included a total of 335 patients with 171 randomized to bendamustine alone and 164 to GAZYVA in combination with bendamustine. With an overall median observation time of 52.2 months (range: 0-100.9 months), there were 66 deaths (40.2%) in the GAZYVA arm and 85 deaths (51.3%) in the bendamustine-alone arm (OS HR = 0.71, 95% CI: 0.51, 0.98). The Kaplan-Meier curve for OS is presented in Figure 4.

Figure 4
Kaplan-Meier Curve of Overall Survival in Patients with FL



GALLIUM

The efficacy of GAZYVA was evaluated in GALLIUM (NCT01332968), a multicenter, open-label, randomized study that included 1202 patients with previously untreated, stage II bulky, III or IV FL. Patients were randomized 1:1 to receive either GAZYVA (n = 601) or rituximab product (n = 601) in combination with chemotherapy (CHOP, CVP, or bendamustine) for 6–8 cycles. Patients were stratified by chemotherapy (selected by each site; all patients at that site received the chosen chemotherapy regimen), FLIPI (Follicular Lymphoma International Prognostic Index) risk group and geographic region. Patients with at least PR to combination therapy received monotherapy with GAZYVA (1,000 mg) or rituximab product every two months until disease progression or for a maximum of two years. The study excluded patients with follicular lymphoma grade 3b or transformed disease; patients having an ANC < 1500 / μ L, platelets < 75,000 / μ L, or CLCr < 40 mL/min; and patients with hepatic transaminases > 2.5 x upper limit of normal unless attributable to lymphoma.

GAZYVA was given by intravenous infusion as a flat dose of 1,000 mg on Days 1, 8 and 15 of Cycle 1 and Day 1 of subsequent treatment cycles.

GAZYVA and bendamustine were given in six 28-day cycles. Bendamustine was administered at 90 mg/m²/day on Days 1 and 2 of each cycle, with prednisone 100 mg orally or equivalent on Day 1 of Cycle 1.

GAZYVA and CHOP were given in six 21-day cycles. Subsequently, two additional cycles of GAZYVA were given for a total of 8 GAZYVA cycles. CHOP consisted of cyclophosphamide 750 mg/m² intravenously, doxorubicin 50 mg/m², and vincristine 1.4 mg/m² (maximum dose, 2 mg) on Day 1 and prednisone 100 mg orally on Days 1-5.

GAZYVA and CVP were given in eight 21-day cycles. CVP consisted of cyclophosphamide 750 mg/m² intravenously and vincristine 1.4 mg/m² (maximum dose, 2 mg) on Day 1 and prednisone 100 mg orally on Days 1-5.

Patients had a median age of 59 years, 81% were White and 53% were female; 7% had Stage II, 35% had Stage III, and 56% had Stage IV disease, with 44% having bulky disease (≥ 7 cm) overall; 79% had a FLIPI score of > 2 ; and 97% had an ECOG performance status of 0–1. The chemotherapy was bendamustine in 57%, CHOP in 33%, and CVP in 10% of patients.

Efficacy was based on PFS per IRC, with a median observation time of 38 months. Upon interim analysis, the risk of progression or death was significantly reduced in the GAZYVA containing arm compared to the rituximab product containing arm (Table 14). Kaplan-Meier curves for PFS are shown in Figure 5. Overall response and complete remission rates were similar.

Table 14 Efficacy in Previously Untreated Follicular Lymphoma (GALLIUM)

Endpoint per IRC	GAZYVA + chemotherapy followed by GAZYVA monotherapy n = 601	Rituximab product + chemotherapy followed by rituximab product monotherapy n = 601
Progression-Free Survival ^a Number of events (%)	108 (18%) HR = 0.72 [95% CI: 0.56, 0.93], p-value = 0.0118 ^b	141 (23%)
Overall Response Rate ^c	91%	88%
Complete Remission Rate ^c	28%	27%


^a Investigator-assessed PFS was consistent with data from independent review.

^b Stratified log-rank test

^c After completion of combination therapy. Assessed by CT without positron emission tomography.

Figure 5

Kaplan-Meier Curves of Progression Free Survival in Patients with Previously Untreated FL



CI, confidence interval; G-chemo, obinutuzumab plus chemotherapy; HR, hazard ratio; NR, not reached; PFS, progression-free survival; R-chemo, rituximab plus chemotherapy

16 HOW SUPPLIED/STORAGE AND HANDLING

GAZYVA is a clear, colorless to slightly brown, preservative-free solution for intravenous use supplied as 1,000 mg/40 mL (25 mg/mL) in single-dose vials (NDC 50242-070-01).

Store at 2°C to 8°C (36°F to 46°F). Do not use beyond expiration date stamped on carton. Protect from light. DO NOT FREEZE. DO NOT SHAKE.

17 PATIENT COUNSELING INFORMATION

Advise patients to seek immediate medical attention for any of the following:

- Signs and symptoms of infusion-related reactions including dizziness, nausea, chills, fever, vomiting, diarrhea, breathing problems, or chest pain [see *Warnings and Precautions (5.3) and Adverse Reactions (6.1)*].
- Symptoms of tumor lysis syndrome such as nausea, vomiting, diarrhea, and lethargy [see *Warnings and Precautions (5.5) and Adverse Reactions (6.1)*].
- Signs of infections including fever and cough [see *Warnings and Precautions (5.6) and Adverse Reactions (6.1)*].
- Symptoms of hepatitis including worsening fatigue or yellow discoloration of skin or eyes [see *Warnings and Precautions (5.1)*].
- New or changes in neurological symptoms such as confusion, dizziness or loss of balance, difficulty talking or walking, or vision problems [see *Warnings and Precautions (5.2)*].

Advise patients of the need for:

- Periodic monitoring of blood counts [*see Warnings and Precautions (5.7 and 5.8) and Adverse Reactions (6.1)*].
- Avoid vaccinations with live viral vaccines [*see Warnings and Precautions (5.9)*].
- Patients with a history of hepatitis B infection (based on the blood test) should be monitored and sometimes treated for their hepatitis [*see Warnings and Precautions (5.1)*].

Advise pregnant women of potential fetal B-cell depletion. Advise females of reproductive potential to inform their healthcare provider of a known or suspected pregnancy [*see Warnings and Precautions (5.10), Use in Specific Populations (8.1)*].

Advise females of reproductive potential to use effective contraception during treatment with GAZYVA and for at least 6 months after the last dose [*see Use in Specific Populations (8.3)*].

Advise women not to breastfeed during treatment with GAZYVA and for at least 6 months after the last dose [*see Use in Specific Populations (8.2)*].

GAZYVA® (obinutuzumab)

Manufactured by:

Genentech, Inc.

A Member of the Roche Group

South San Francisco, CA 94080-4990

U.S. License No. 1048

GAZYVA is a registered trademark of Genentech, Inc.

© 2020 Genentech, Inc.