

NOW APPROVED

SET A HIGHER BAR VS PLACEBO + ST IN LUPUS NEPHRITIS¹

GAZYVA + ST showed superior CRR vs placebo + ST
46.4% of patients achieved CRR vs 33.1%, with a treatment difference of 13.4% (95% CI, 2.0, 24.8; $P=0.0232$)^{1*†}

Actor portrayal.

EULAR 2025 Guidelines recommend early use of GAZYVA²

For patients with active lupus nephritis, especially those with poor prognostic factors, GAZYVA is a recommended option in combination with standard therapy.

Indication

GAZYVA[®] (obinutuzumab) is indicated for the treatment of adult patients with active lupus nephritis (LN) who are receiving standard therapy.

Important Safety Information

BOXED WARNINGS: HEPATITIS B VIRUS REACTIVATION AND PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY

- Hepatitis B Virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients receiving CD20-directed cytolytic antibodies, including GAZYVA. Screen all patients for HBV infection before treatment initiation. Monitor HBV-positive patients during and after treatment with GAZYVA. Discontinue GAZYVA and concomitant medications in the event of HBV reactivation
- Progressive Multifocal Leukoencephalopathy (PML), including fatal PML, can occur in patients receiving GAZYVA

Contraindications

- GAZYVA is contraindicated in patients with known hypersensitivity reactions (eg, anaphylaxis) to obinutuzumab or to any of the excipients, or serum sickness with prior obinutuzumab use

Please see additional Important Safety Information throughout and the full [**Prescribing Information**](#), including **BOXED WARNINGS**.

GAZYVA[®]
obinutuzumab
injection | 1,000mg/40mL

*CRR at Week 76: defined as UPCR <0.5 g/g; an eGFR ≥85% of baseline, with no occurrence of rescue therapy, treatment failure, death, or early study withdrawal through Week 76.¹

[†]Standard therapy (ST) included MMF 2 g/d to 2.5 g/d and a tapering course of corticosteroids to a target dose of 5 mg/d.¹

CI=confidence interval; CRR=complete renal response; eGFR=estimated glomerular filtration rate; EULAR=European Alliance of Associations for Rheumatology; MMF=mycophenolate mofetil.



About LN/MOA

Study Design

Efficacy

Safety

Dosing/Guidelines

Access/Support

Summary

Lupus nephritis carries serious risks that demand urgent attention³⁻⁸



Lupus nephritis does not always show clear warning signs⁴

- Lupus nephritis can silently affect the kidneys, quickly escalating from mild subclinical nephritis to severe, rapidly progressive glomerulonephritis⁴
- Up to 28% of patients with lupus nephritis go on to develop end-stage renal disease, highlighting the substantial risk for disease progression⁵



Renal outcomes can be especially poor for patients who:

Do not achieve CRR*

- Retrospective observational studies suggest that patients who do not achieve CRR may face a higher risk of renal failure^{3,6,7}
- Opportunity remains to improve CRR rates beyond standard therapy alone²

Have suboptimal treatment adherence

- ~65% of patients with lupus nephritis are not adherent to treatment, increasing their risk of renal flares⁸
- Common reasons for lack of adherence include depression, low socioeconomic status, rural residence, and polypharmacy, according to a systematic review of 11 studies⁹



Current ACR, KDIGO, and EULAR guidelines recommend prompt use of triple therapy and IV administration to address treatment gaps^{2,10,11†‡}

- ACR 2024 Guidelines conditionally recommend triple therapy as a preferred initial treatment for patients with active, newly diagnosed, or flaring disease^{‡§}
- KDIGO 2024 Guidelines recommend triple therapy as an initial treatment for patients with active class III or IV ± V lupus nephritis[‡]
 - KDIGO guidelines also state that physicians may choose an IV regimen if suboptimal adherence is anticipated[‡]
- EULAR 2025 Guidelines recommend early use of combination therapy for patients with active lupus nephritis, especially those with poor prognostic factors

*CRR was defined in the retrospective studies as achieving eGFR within 10% of baseline or ≥90 mL/min/1.73 m² and low proteinuria (typically UPCR <0.5 g/g or 24-hr protein <0.5 g/d); one study applied modified ALMS and BLISS-LN criteria, requiring normal serum creatinine or estimated creatinine clearance and proteinuria ≤0.5 g/d, excluding urinary sediment, all within 2 years.^{3,6,7}

†Triple therapy is defined as immunosuppressants, steroids, and advanced therapy.^{2,10,11}

‡As of September 2025, ACR and KDIGO guidelines do not include obinutuzumab.^{10,11}

§For the full 2024 ACR Lupus Nephritis Guidelines, please visit rheumatology.org/lupus-guideline.¹⁰

ACR=American College of Rheumatology; ALMS=Aspreva Lupus Management Study; BLISS-LN=Belimumab International Study in Lupus Nephritis; CRR=complete renal response; eGFR=estimated glomerular filtration rate; EULAR=European Alliance of Associations for Rheumatology; KDIGO=Kidney Disease: Improving Global Outcomes; UPCR=urine protein-to-creatinine ratio.


GAZYVA is the first and only FDA-approved B-cell depleter in lupus nephritis¹

B cells are a key driver of irreversible kidney damage in lupus nephritis¹²⁻¹⁴



GAZYVA is a type II, humanized, anti-CD20 monoclonal antibody that targets B cells^{1,15}

How the distinct design of GAZYVA (obinutuzumab, type II) compares to rituximab (type I) based on preclinical data^{1,15,16}:



Nearly all patients in the REGENCY study experienced B-cell depletion by Week 4, sustained through Week 76.^{11¶}

[¶]Defined as B-cell count <10 cells/μL. B-cell depletion was observed in 99% of patients by Week 4 and remained below this threshold in 95% of patients at Week 76.¹

[¶]B-cell depletion has not been shown to be directly correlated to clinical response.¹

ADCC=antibody-dependent cellular cytotoxicity; CD20=cluster of differentiation 20; Fc=crystallizable fragment.

Important Safety Information (cont'd)

Warnings and Precautions

Hepatitis B Virus Reactivation

- GAZYVA can cause hepatitis B virus (HBV) reactivation. HBV reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients treated with anti-CD20 antibodies, including GAZYVA. HBV reactivation has been reported in patients who are hepatitis B surface antigen (HBsAg) positive and in patients who are HBsAg negative but are hepatitis B core antibody (anti-HBc) positive. Reactivation has also occurred in patients who appear to have resolved hepatitis B infection (ie, HBsAg negative, anti-HBc positive, and hepatitis B surface antibody [anti-HBs] positive)

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.



REGENCY evaluated the efficacy and safety of GAZYVA + ST* vs placebo + ST¹

REGENCY was a Phase III, randomized, double-blind, placebo-controlled, multicenter study^{1,18}

Eligibility criteria

- Diagnosis of active ISN/RPS class III or IV ± class V lupus nephritis[‡]
- UPCR ≥1 g/g on a 24-hr collection
- ≥1 dose of pulse methylprednisolone IV (≥250 mg) or equivalent, 6 months prior to or during screening

Randomized 1:1 (N=271).

Primary analysis at Week 76

GAZYVA + ST (n=135)

1000 mg IV GAZYVA on Weeks 0, 2, 24, 26, 50,[‡] and 52.
[‡]Patients received either placebo or GAZYVA at Week 50.

Patients received treatment every 6 months; inadequate responders entered open-label use. All were followed for at least 12 months after Week 76 treatment.

Placebo + ST (n=136)

Placebo on Weeks 0, 2, 24, 26, 50, and 52.

Primary endpoint¹

- Complete renal response (CRR)

Select key secondary endpoints^{1,19}

- CRR with successful prednisone taper between Weeks 64 and 76 (≤7.5 mg/d)
- Change in eGFR from baseline
- Proportion of patients with renal-related event or death

Exploratory endpoints^{1,18}

- Components of CRR
- CRR subgroup analysis
- Time to renal-related event or death
- Time to lupus nephritis flares between Weeks 24 and 76

*Standard therapy (ST) included mycophenolate mofetil (MMF) 2 g/d to 2.5 g/d and a tapering course of corticosteroids to a target dose of 5 mg/d.¹
[‡]Renal biopsy performed within 6 months before screening. Classified according to the ISN/RPS 2003 classification system for lupus nephritis.¹
eGFR=estimated glomerular filtration rate; ISN/RPS=International Society of Nephrology/Renal Pathology Society; SD=standard deviation; UPCR=urine protein-to-creatinine ratio.

Important Safety Information (cont'd)
Warnings and Precautions (cont'd)
Hepatitis B Virus Reactivation (cont'd)

- HBV reactivation is defined as an abrupt increase in HBV replication manifesting as a rapid increase in serum HBV DNA level, or detection of HBsAg in a person who was previously HBsAg negative and anti-HBc positive. Reactivation of HBV replication is often followed by hepatitis, ie, increase in transaminase levels and in severe cases, increase in bilirubin levels, liver failure, and death
- Screen all patients for HBV infection by measuring HBsAg and anti-HBc before initiating treatment with GAZYVA. For patients who show evidence of hepatitis B infection (HBsAg positive [regardless of antibody status] or HBsAg negative but anti-HBc positive), consult healthcare providers with expertise in managing hepatitis B regarding monitoring and consideration for HBV antiviral therapy
- Monitor patients with evidence of current or prior HBV infection for clinical and laboratory signs of hepatitis or HBV reactivation during and for several months following treatment with GAZYVA

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.

The REGENCY study population was well-represented across key clinical characteristics^{1,18}

A diverse and inclusive study population^{1,18}

Select demographics and baseline characteristics combining both arms (N=271)

Age

Median (years)	31
----------------	----

Sex

Female	85%
--------	-----

Ethnicity/race[‡]

Hispanic or Latino	58%
White	48%
American Indian or Alaska Native	19%
Black or African American	15%
Asian	6%

Kidney biopsy class

Class III	39%
Class IV	61%
Concomitant class V	31%

eGFR at baseline (mL/min/1.73 m²)

Mean	102.3 (SD: ±30.8)
<90	29%
≥90	71%

24-hr UPCR at baseline (mg/mg)

Mean	3.3 (SD: ±2.9)
<3	58%
≥3	42%

Disease history


Newly diagnosed	42%
Prior history	58%
Median lupus nephritis duration if prior history (months)	37 (range: 0.4-330.4)
No prior advanced therapy [§]	93%

[‡]Ethnicity and race were not stated or were unknown for some patients. Patients could report multiple races.¹⁹
[§]Prior advanced therapies had to be discontinued >2 months before or during study screening.¹



Nearly half of patients treated with GAZYVA + ST* achieved CRR¹

Primary endpoint
Superior CRR at Week 76^{1†‡}



Components of CRR in REGENCY ^{1,18}	Exploratory endpoint	
	GAZYVA + ST (n=135)	Placebo + ST (n=136)
UPCR <0.5 g/g	47.4%	36.0%
eGFR ≥85% of baseline [§]	83.7%	75.7%
No occurrence of intercurrent events	88.9%	75.0%

Exploratory analyses were not type I error-controlled; therefore, no conclusions can be drawn.

*Standard therapy (ST) included mycophenolate mofetil (MMF) 2 g/d to 2.5 g/d and a tapering course of corticosteroids to a target dose of 5 mg/d.¹
¹CRR at Week 76: defined as UPCR <0.5 g/g; an eGFR ≥85% of baseline, with no occurrence of rescue therapy, treatment failure, death, or early study withdrawal through Week 76.¹
[†]Analyzed using the Cochran-Mantel-Haenszel test, adjusted for race and region. Missing data were handled using the multiple-imputation method (4 patients in the GAZYVA + ST arm; 1 patient in the placebo + ST arm).^{1,19}
[§]As calculated using the 2009 CKD-EPI equation.¹
^{||}Rescue therapy, treatment failure, death, or early study withdrawal.¹


Important Safety Information (cont'd) Warnings and Precautions (cont'd) Hepatitis B Virus Reactivation (cont'd)

- In patients who develop reactivation of HBV while receiving GAZYVA, immediately discontinue GAZYVA and any concomitant chemotherapy and institute appropriate treatment. Resumption of GAZYVA in patients whose HBV reactivation resolves should be discussed with healthcare providers with expertise in managing hepatitis B. Insufficient data exist regarding the safety of resuming GAZYVA in patients who develop HBV reactivation

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.

GAZYVA + ST achieved CRR while also reducing steroid burden¹

Secondary endpoint
CRR at Week 76 with successful steroid taper vs placebo + ST^{1‡}



REGENCY steroid-tapering protocol

- All patients received oral prednisone on a standardized tapering schedule of 0.5 mg/kg, with a target of reaching 7.5 mg/d by Week 12 and 5 mg/d by Week 24, maintained through Week 80^{1,19}
- Successful steroid tapering was defined as maintaining ≤7.5 mg/d between Weeks 64 and 76¹

CI=confidence interval; CKD-EPI=Chronic Kidney Disease Epidemiology Collaboration; CRR=complete renal response; eGFR=estimated glomerular filtration rate; UPCR=urine protein-to-creatinine ratio.

Important Safety Information (cont'd) Warnings and Precautions (cont'd) Progressive Multifocal Leukoencephalopathy (PML)

- John Cunningham (JC) virus infection resulting in PML, which can be fatal, occurred in patients treated with GAZYVA in chronic lymphocytic leukemia (CLL) and non-Hodgkin lymphoma (NHL). Consider the diagnosis of PML in any patient presenting with new onset or changes to, preexisting neurologic manifestations. Evaluation of PML includes, but is not limited to, consultation with a neurologist, brain MRI, and lumbar puncture. Discontinue GAZYVA therapy and consider discontinuation or reduction of any concomitant chemotherapy or immunosuppressive therapy in patients who develop PML

Infusion-Related Reactions

- GAZYVA can cause severe and life-threatening infusion-related reactions (IRRs)
- In patients with LN, IRRs occurred predominantly during infusion of the first 1,000 mg. IRRs were generally mild to moderate and could be managed by slowing or temporarily halting the infusion. Severe and life-threatening IRRs requiring symptomatic treatment were also reported. The most common IRR signs/symptoms reported in the REGENCY study included headache, nausea, and vomiting. In the NOBILITY study, the most common IRR symptoms were pyrexia and tachycardia
- Premedicate patients with acetaminophen, an antihistamine, and a glucocorticoid. Closely monitor patients during the entire infusion. Reduce infusion rate, interrupt infusion, or permanently discontinue GAZYVA for IRRs based on severity. Institute medical management (eg, glucocorticoids, epinephrine, bronchodilators, and/or oxygen) for IRRs as needed



CRR was evaluated across prespecified subgroups^{1,19}

Exploratory endpoint
CRR at Week 76 across baseline disease characteristics

Subgroup	Number of patients GAZYVA + ST*	Placebo + ST	Patients with CRR GAZYVA + ST	Placebo + ST	Treatment difference Percentage points (95% CI)
All patients	135	136	46%	33%	13.4 (2.0, 24.8)
History of lupus nephritis at baseline					
Previously diagnosed	81	76	43%	38%	6.0 (-9.2, 21.3)
Newly diagnosed	54	60	52%	27%	23.8 (5.7, 39.8)
Lupus nephritis class at baseline					
Class III	56	51	50%	41%	7.1 (-11.5, 25.6)
Class IV	79	85	44%	28%	15.2 (0.8, 29.5)
Concomitant class V disease at baseline					
No	88	98	45%	37%	8.8 (-5.2, 22.8)
Yes	47	38	49%	24%	25.5 (6.3, 44.7)
24-hr UPCR at baseline					
<3	82	74	52%	46%	7.8 (-7.7, 23.2)
≥3	52	62	37%	18%	17.8 (1.6, 33.9)

Subgroup findings should be interpreted with caution due to small sample sizes and overlapping subgroups.

CI=confidence interval; eGFR=estimated glomerular filtration rate; SD=standard deviation; SE=standard error; UPCR=urine protein-to-creatinine ratio.

Important Safety Information (cont'd) Warnings and Precautions (cont'd) Infusion-Related Reactions (cont'd)

- For patients with preexisting cardiac or pulmonary conditions, monitor more frequently throughout the infusion and the post-infusion period, since they may be at greater risk of experiencing more severe reactions. Hypotension may occur as part of the IRR to GAZYVA. Consider withholding antihypertensive treatments for 12 hours prior to, during each GAZYVA infusion, and for the first hour after administration until blood pressure is stable. For patients at increased risk of hypertensive crisis, consider the benefits versus the risks of withholding their antihypertensive medication

Hypersensitivity Reactions Including Serum Sickness

- Hypersensitivity reactions have been reported in patients treated with GAZYVA. Signs of immediate-onset hypersensitivity included dyspnea, bronchospasm, hypotension, urticaria, and tachycardia. Late-onset hypersensitivity diagnosed as serum sickness has also been reported, with symptoms that include chest pain, diffuse arthralgia, and fever. Hypersensitivity reactions may be difficult to clinically distinguish from IRRs. However, hypersensitivity very rarely occurs with the first infusion and, when observed, often occurs after previous exposure

Difference in eGFR from baseline was evaluated^{18,19}

Secondary endpoint
Change in eGFR from baseline through Week 76^{18,19†}

- GAZYVA + ST showed a numerical increase in eGFR



- Baseline eGFR levels: 102.8 mL/min/1.73 m² (SD: 29.3) in the GAZYVA + ST arm compared with 101.9 mL/min/1.73 m² (SD: 32.2) for placebo + ST^{18,19}

*Standard therapy (ST) included mycophenolate mofetil (MMF) 2 g/d to 2.5 g/d and a tapering course of corticosteroids to a target dose of 5 mg/d.¹
†Missing data were handled using the multiple-imputation method (8 patients in the GAZYVA + ST arm; 12 patients in the placebo + ST arm).¹⁹

Important Safety Information (cont'd) Warnings and Precautions (cont'd) Hypersensitivity Reactions Including Serum Sickness (cont'd)

- If a hypersensitivity reaction is suspected during or after an infusion, stop the infusion and permanently discontinue treatment. GAZYVA is contraindicated in patients with known hypersensitivity reactions to GAZYVA, including serum sickness with prior GAZYVA use

Serious, Including Fatal, Infections

- Fatal and serious bacterial, fungal, and new or reactivated viral infections can occur during and following GAZYVA therapy
- In the pooled double-blind periods of REGENCY (Week 76) and NOBILITY (Week 52), the incidence of Grade 3-5 infections was 11% (22/200) in patients treated with GAZYVA and standard therapy compared to 9% (18/193) in patients receiving placebo and standard therapy, corresponding to an exposure-adjusted incidence rate (EAIR) of 8.9 and 7.9 per 100 patient years, respectively. The incidence of fatal infections was 1% (2/200) in patients treated with GAZYVA and 0.5% (1/193) in patients receiving placebo, corresponding to an EAIR of 0.8 and 0.4 per 100 patient years, respectively
- In 40 patients who crossed-over from placebo to GAZYVA and standard therapy at Week 76 in the REGENCY study, and patients who continued treatment with GAZYVA and standard therapy, including additional treatment after Week 76, the EAIR of Grade 3-5 infections for the GAZYVA arm was 9.0 per 100 patient years while the EAIR of fatal infections for the GAZYVA arm was 1.8 per 100 patient years

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.

GAZYVA
obinutuzumab
injection 1,000mg/40mL

Fewer patients experienced a renal-related event or death with GAZYVA + ST* vs placebo + ST¹


Secondary endpoint

Proportion of patients with renal-related event or death through Week 76

- 17.8% (n=24/135) with GAZYVA + ST vs 33.8% (n=46/136) with placebo + ST
- Due to the hierarchical design, this analysis was not tested for statistical significance; therefore, no conclusions can be drawn

Exploratory endpoint

Time to renal-related event or death



Exploratory analyses were not type I error-controlled; therefore, no conclusions can be drawn.

Components of renal-related event or death in REGENCY	GAZYVA + ST (n=135)	Placebo + ST (n=136)
Treatment failure	4.4%	18.4%
End-stage renal disease (ESRD)	0%	1.5%
Clinically significant worsening in UPCR and/or eGFR from Week 24 [†]	3.7%	16.2%
Rescue therapy (except corticosteroid-only rescue)	3.7%	16.2%
Worsening proteinuria [‡]	12.6%	13.2%
Worsening eGFR [§]	5.2%	14.7%
Death	2.2%	0.7%

- 3.0% of patients had a doubling of serum creatinine from baseline with GAZYVA + ST vs 5.9% with placebo + ST

*Standard therapy (ST) included mycophenolate mofetil (MMF) 2 g/d to 2.5 g/d and a tapering course of corticosteroids to a target dose of 5 mg/d.¹
[†]Clinically significant, sustained worsening in UPCR and/or eGFR that leads the investigator to conclude the patient has failed the randomized treatment period.¹
[‡]Worsening proteinuria was prospectively defined as a confirmed ≥50% increase in UPCR to a value ≥3 g/g.¹
[§]Worsening eGFR was prospectively defined as confirmed ≥30% decrease in eGFR to a value <60 mL/min/1.73 m².¹

Important Safety Information (cont'd)

Warnings and Precautions (cont'd)

Serious, Including Fatal, Infections (cont'd)


- Do not administer GAZYVA to patients with an active infection. Patients with a history of recurring or chronic infections may be at increased risk of infection. In patients who develop a serious infection while receiving GAZYVA, immediately discontinue GAZYVA and institute appropriate treatment. Consider the risk and benefit of resuming treatment with GAZYVA following resolution of serious infections

Time to renal flare was evaluated in a prespecified exploratory analysis¹⁸

Exploratory endpoint

Proportion of patients with lupus nephritis flares from Week 24 through Week 76¹⁸

- 11.1% (n=15/135) with GAZYVA + ST vs 23.5% (n=32/136) with placebo + ST



Exploratory analyses were not type I error-controlled; therefore, no conclusions can be drawn.

Renal flares, as defined in REGENCY¹⁹

Patients exhibited one of the following:

- eGFR decrease to >20% compared with Week 24 in patients with UPCR >1 g/g and/or cellular casts
- UPCR increase to:
 - >1 g/g if Week 24 UPCR was <0.2 g/g
 - >2 g/g if Week 24 UPCR was 0.2 g/g to 1 g/g
 - Doubling, if Week 24 UPCR was >1 g/g
- Receipt of rescue therapy, except for corticosteroid-only rescue

CI=confidence interval; eGFR=estimated glomerular filtration rate; HR=hazard ratio; UPCR=urine protein-to-creatinine ratio.

Important Safety Information (cont'd)

Warnings and Precautions (cont'd)

Neutropenia

- Severe and life-threatening neutropenia, including febrile neutropenia, has been reported during treatment with GAZYVA. Monitor patients with Grade 3 to 4 neutropenia frequently with regular laboratory tests until resolution. Anticipate, evaluate, and treat any symptoms or signs of developing infection. Consider dose delays for Grade 3 or 4 neutropenia. Consider administration of granulocyte colony-stimulating factors (GCSFs) in patients with Grade 3 or 4 neutropenia
- Neutropenia can also be of late onset (occurring more than 28 days after completion of treatment) and/or prolonged (lasting longer than 28 days)
- Patients with severe and long-lasting (>1 week) neutropenia are strongly recommended to receive antimicrobial prophylaxis until resolution of neutropenia to Grade 1 or 2. Consider antiviral and antifungal prophylaxis

Thrombocytopenia

- Severe and life-threatening thrombocytopenia has been reported during treatment with GAZYVA in combination with chemotherapy. Fatal hemorrhagic events have been reported in patients with CLL treated with GAZYVA in combination with chemotherapy
- Monitor patients frequently for thrombocytopenia and hemorrhagic events, and if clinically indicated, evaluate laboratory coagulation parameters. In patients with Grade 3 or 4 thrombocytopenia, monitor platelet counts more frequently until resolution and consider dose delays of GAZYVA and chemotherapy or dose reductions of chemotherapy. Transfusion of blood products (ie, platelet transfusion) may be necessary. Consider withholding concomitant medications that may increase bleeding risk (platelet inhibitors or anticoagulants), especially during the first cycle

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.



GAZYVA has a well-demonstrated safety profile^{1,20}

Pooled ARs up to Week 76 (≥2% greater in the GAZYVA + ST* arm) across 2 double-blind, placebo-controlled studies: Phase III REGENCY and Phase II NOBILITY^{††}

Adverse reactions	GAZYVA + ST (n=200)		Placebo + ST (n=193)	
	All grades (%)	Grades 3-5 (%)	All grades (%)	Grades 3-5 (%)
Infections and infestations				
Upper respiratory tract infection	29	0	24	0
COVID-19	23	5	16	0.5
Urinary tract infection	21	3	18	4
Bronchitis	14	0	8	0.5
Pneumonia	10	2	6	2
Herpes simplex	3	0	0	0
Injury, poisoning, and procedural complications				
Infusion-related reaction	14	2	10	0.5
Blood and lymphatic system disorders				
Neutropenia	14	7	6	0.5

- The most common serious ARs in the GAZYVA + ST arm were COVID-19 (5.5%), pneumonia (4.5%), neutropenia (3.5%), urinary tract infections (2.5%), and infusion-related reactions (0.5%). Two fatal ARs of COVID-19 were reported in the GAZYVA + ST arm[†]

Most ARs were grade 1-2.[†]

*Standard therapy (ST): mycophenolate mofetil (MMF) and corticosteroids for REGENCY, and MMF/MPA and corticosteroids for NOBILITY.[†]
[†]NOBILITY was a Phase II, double-blind, placebo-controlled study that included 64 patients treated with GAZYVA plus standard therapy consisting of MMF/MPA and corticosteroids.^{1,20}

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.

Additional safety considerations with GAZYVA + ST

Safety highlights from REGENCY and NOBILITY

IRRs[†]

- IRRs in both arms were predominantly grade 1-2 and occurred during or after the first infusion
- All grade 3-4 events occurred during or after either the first or second infusion
- The incidence and severity of IRRs decreased with subsequent infusions
- No grade 5 IRRs were reported

Neutropenia[†]

- Grade 3-4 neutropenia was reported in 7% of patients treated with GAZYVA + ST vs 0.5% of patients receiving placebo + ST
- Neutropenia and related AEs resolved or improved spontaneously or with use of granulocyte colony-stimulating factors in 96% of patients

Infections

- Infections were reported in 72% of patients in the GAZYVA + ST arm vs 62% of patients in the placebo + ST arm[†]
- Grade 3-5 infections events were reported in 11% of patients in the GAZYVA + ST arm vs 10% of patients in the placebo + ST arm[†]
- Fatal infections were reported in 1% of patients in the GAZYVA + ST arm vs 0.5% of patients in the placebo + ST arm^{1,18}

REGENCY safety

Infections and the impact of COVID-19

- REGENCY began in July 2020, during the peak of the global COVID-19 pandemic¹⁹
- Excluding COVID-19 events, the incidence of serious infections decreased from 17% to 11% in the GAZYVA + ST arm, with no change in the placebo + ST arm (8%)¹⁹
- No COVID-19-related serious ARs occurred during the latter half of the study when vaccines were widely available; 57% of patients on GAZYVA received COVID-19 vaccines during the study^{18,19}

Discontinuation rates¹⁹

- ARs leading to discontinuation occurred in 7% of patients receiving GAZYVA + ST and 4% of patients with placebo + ST

AE=adverse event; AR=adverse reaction; IRR=infusion-related reaction; MPA=mycophenolic acid.



GAZYVA offers twice-yearly IV dosing¹


After 4 initial doses, patients receive only 1 dose once every 6 months¹



HCP administration may enable closer monitoring of patient adherence.^{21,22}

HCP=healthcare professional; IRR=infusion-related reaction.

EULAR 2025 Guidelines recommend early use of GAZYVA²



Actor portrayal.

For patients with active lupus nephritis, especially those with poor prognostic factors, GAZYVA is a recommended option in combination with standard therapy.²

Important Safety Information (cont'd) Warnings and Precautions (cont'd) Disseminated Intravascular Coagulation (DIC)

- Fatal and severe DIC has been reported in patients receiving GAZYVA for CLL and NHL. The majority of DIC cases have involved changes in platelets and laboratory coagulation parameters following the first infusion, with spontaneous resolution usually occurring by Day 8. In some cases, DIC was associated with IRRs
- In patients with suspected DIC, evaluate potential causes and monitor coagulation parameters, platelet counts, and for signs and symptoms of bleeding or thrombosis. Manage according to standard guidelines for DIC. Supportive care, including transfusion of blood products and other medical management, may be necessary

Immunization

- The safety and efficacy of immunization with live or attenuated viral vaccines during or following GAZYVA therapy have not been studied. Immunization with live virus vaccines is not recommended during treatment with GAZYVA and until B-cell recovery

Embryo-Fetal Toxicity

- Based on its mechanism of action and findings in animals, GAZYVA can cause B-cell depletion in infants exposed to obinutuzumab in utero. Advise pregnant women of the potential risk to the fetus. Mothers who have been exposed to GAZYVA during pregnancy should discuss the safety and timing of live virus vaccinations for their infants with their child's healthcare providers. Advise females of reproductive potential to use effective contraception while receiving GAZYVA and for 6 months after the last dose

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.

GAZYVA
obinutuzumab
injection 1,000mg/40mL



Access GAZYVA with confidence

Billing and coding

A permanent J-code is already in place.

This coding information may assist you as you complete the payer forms.

J9301
Injection, obinutuzumab, 10 mg

Access and reimbursement



Genentech Immunology Access Solutions

Offers a range of access and reimbursement support for your patients and practice*:

- Benefits investigations
- Prior authorization resources
- Resources for denials and appeals

Learn more and enroll in Access Solutions today.



Visit genentech-access.com/hcp for more information



Call (866) 681-3261
Support is available Monday through Friday, 9 AM to 8 PM ET

*The completion and submission of coverage- or reimbursement-related documentation are the responsibility of the patient and healthcare provider. Genentech makes no representation or guarantee concerning coverage or reimbursement for any service or item.

Please see Important Safety Information throughout and the full [Prescribing Information](#), including **BOXED WARNINGS**.

Access support is available to help your patients start and stay on GAZYVA

Financial assistance



GAZYVA Immunology Co-pay Program

Provides financial assistance to eligible commercially insured patients who may pay **as little as \$0 co-pay.**[†]

- Up to \$15,000 for GAZYVA drug out-of-pocket (OOP) costs per calendar year
- Up to \$2,000 for GAZYVA administration OOP costs per calendar year

Live support from Case Managers



Genentech Immunology Access Solutions

Provides referrals to financial assistance options throughout your patients' treatment.[‡]

[†]The Co-pay Program ("Program") is valid ONLY for patients with commercial (private or non-governmental) insurance who have a valid prescription for a Food and Drug Administration (FDA)-approved indication of a Genentech medicine. The Program is not available to patients whose prescriptions are reimbursed under any federal, state, or government-funded insurance programs (included but not limited to Medicare, Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs Programs) or where prohibited by law or by the patient's health insurance provider. If at any time a patient begins receiving prescription drug coverage under any such federal, state or government-funded healthcare programs, the patient will no longer be eligible for the Program.

Under the Program, the patient may be required to pay a co-pay. The final amount owed by a patient may be as little as \$0 for the Genentech medicine (see Program specific details available at the Program website). The total patient out-of-pocket cost is dependent on the patient's health insurance plan. The Program assists with the cost of the Genentech medicine only. It does not assist with the cost of other medicines, procedures or office visit fees. After reaching the maximum annual Program benefit amount, the patient will be responsible for all remaining out-of-pocket expenses. The Program benefit amount cannot exceed the patient's out-of-pocket expenses for the Genentech medicine. All participants are responsible for reporting the receipt of all Program benefits as required by any insurer or by law. The Program is only valid in the United States and U.S. Territories, is void where prohibited by law and shall follow state restrictions in relation to AB-rated generic equivalents (eg, MA, CA) where applicable. No party may seek reimbursement for all or any part of the benefit received through the Program. The value of the Program is intended exclusively for the benefit of the patient. The funds made available through the Program may only be used to reduce the out-of-pocket costs for the patient enrolled in the Program. The Program is not intended for the benefit of third parties, including without limitation third-party payers, pharmacy benefit managers, or their agents. If Genentech determines that a third party has implemented a program that adjusts patient cost-sharing obligations based on the availability of support under the Program and/or excludes the assistance provided under the Program from counting towards the patient's deductible or out-of-pocket cost limitations, Genentech may impose a per fill cap on the cost-sharing assistance available under the Program. Submission of true and accurate information is a requirement for eligibility and Genentech reserves the right to disqualify patients who do not comply with Genentech Program Terms and Conditions. Genentech reserves the right to rescind, revoke, or amend the Program without notice at any time.

Additional terms and conditions apply. Please visit the Co-pay Program website for the full list of Terms and Conditions.

[‡]Genentech provides coverage and reimbursement services to patients to help them understand benefits, coverage, and reimbursement. Genentech provides these services to patients only after a healthcare provider has prescribed a Genentech product.

GAZYVA
obinutuzumab
injection | 1,000mg/40mL



GAZYVA Clinical Education Managers are available to help your patients

Personal support from Day 1 for patients on their GAZYVA journey

The more patients know about their treatment, the more actively involved they can be in their lupus nephritis care. GAZYVA Clinical Education Managers (CEMs) help provide a strong foundation for the first infusion and beyond—with in-person or virtual educational sessions, disease information, and treatment resources, all at no cost.

CEMs can provide your patients:



No-cost in-person or virtual educational sessions



Resources and information about lupus nephritis



Help preparing for the first GAZYVA infusion



One-on-one support from our dedicated team

CEMs do not provide medical advice. If patients have questions about their health, encourage them to contact their healthcare provider.

Encourage your patients to schedule their first session with a CEM by calling 1-866-681-3261.

Important Safety Information (cont'd)

Important Safety Information (cont'd)

Warnings and Precautions (cont'd)

Lactation

- Human IgG is known to be present in human milk. Because of the potential of serious adverse reactions in the breastfed child, advise women not to breastfeed during treatment with GAZYVA and for 6 months after the last dose

Additional Important Safety Information

- The most common adverse reactions (incidence $\geq 5\%$) observed in patients with LN in the GAZYVA arm were upper respiratory tract infection, COVID-19, urinary tract infection, bronchitis, pneumonia, infusion-related reactions, and neutropenia
- The most common serious adverse reactions in the GAZYVA arm were: COVID-19 (5.5%), pneumonia (4.5%), neutropenia (3.5%), urinary tract infections (2.5%), and infusion-related reactions (0.5%). No serious adverse reactions were reported for bronchitis, herpes simplex and upper respiratory tract infections. Two fatal adverse reactions of COVID-19 were reported in the GAZYVA arm

You are encouraged to report side effects to Genentech and the FDA. You may contact Genentech by calling 1-888-835-2555. You may contact the FDA by visiting www.fda.gov/medwatch or calling 1-800-FDA-1088.

Please see the full [Prescribing Information](#) for additional Important Safety Information, including **BOXED WARNINGS**.

References: 1. GAZYVA full Prescribing Information. South San Francisco, CA: Genentech, Inc.; 2025. 2. Fanouriakis A, Kostopoulou M, Anders HJ, et al. EULAR recommendations for the management of systemic lupus erythematosus with kidney involvement: 2025 update. *Ann Rheum Dis*. Published online October 16, 2025. doi:10.1016/j.ard.2025.09.007 3. Urowitz M, Georgiou ME, Touma Z, et al. Renal response status to predict long-term renal survival in patients with lupus nephritis: results from the Toronto Lupus Cohort. *Lupus Sci Med*. 2024;11(2):e001264. doi:10.1136/lupus-2024-001264 4. Anders HJ, Saxena R, Zhao MH, Parodis I, Salmon JE, Mohan C. Lupus nephritis. *Nat Rev Dis Primers*. 2020;6(1):7. doi:10.1038/s41572-019-0141-9 5. Mahajan A, Amelio J, Gairy K, et al. Systemic lupus erythematosus, lupus nephritis and end-stage renal disease: a pragmatic review mapping disease severity and progression. *Lupus*. 2020;29(9):1011-1020. doi:10.1177/0961203320932219 6. Yap DYH, Xu X, Juliao PC, et al. Long-term kidney outcome of lupus nephritis by renal response status. *Kidney Int Rep*. 2024;9(12):3532-3541. doi:10.1016/j.ekir.2024.09.028 7. Davidson JE, Fu Q, Ji B, et al. Renal remission status and longterm renal survival in patients with lupus nephritis: a retrospective cohort analysis. *J Rheumatol*. 2018;45(5):671-677. doi:10.3899/jrheum.161554 8. Ali AY, Abdelaziz TS, Behiry ME. The prevalence and causes of non-adherence to immunosuppressive medications in patients with lupus nephritis flares. *Curr Rheumatol Rev*. 2020;16(3):245-248. doi:10.2174/1573397115666190626111847 9. Mehat P, Atiquzzaman M, Esdaile JM, Aviña-Zubieta A, De Vera MA. Medication nonadherence in systemic lupus erythematosus: a systematic review. *Arthritis Care Res (Hoboken)*. 2017;69(11):1706-1713. doi:10.1002/acr.23191 10. Sammaritano LR, Askanase A, Bermas BL, et al. 2024 American College of Rheumatology (ACR) guideline for the screening, treatment, and management of lupus nephritis. *Arthritis Care Res (Hoboken)*. 2025;77(9):1045-1065. doi:10.1002/acr.25528 11. Kidney Disease: Improving Global Outcomes (KDIGO) Lupus Nephritis Work Group. KDIGO 2024 clinical practice guideline for the management of lupus nephritis. *Kidney Int*. 2024;105(suppl 1):S1-S69. doi:10.1016/j.kint.2023.09.002 12. Gomez Mendez LM, Cascino MD, Garg J, et al. Peripheral blood B cell depletion after rituximab and complete response in lupus nephritis. *Clin J Am Soc Nephrol*. 2018;13(10):1502-1509. doi:10.2215/CJN.01070118. Published correction appears in *Clin J Am Soc Nephrol*. 2019;14(1):111. doi:10.2215/CJN.12621018 13. Lech M, Anders HJ. The pathogenesis of lupus nephritis. *J Am Soc Nephrol*. 2013;24(9):1357-1366. doi:10.1681/ASN.2013010026 14. Thompson JC, Mahajan A, Scott DA, Gairy K. The economic burden of lupus nephritis: a systematic literature review. *Rheumatol Ther*. 2022;9(1):25-47. doi:10.1007/s40744-021-00368-y 15. Herter S, Herting F, Mundigl O, et al. Preclinical activity of the type II CD20 antibody GA101 (obinutuzumab) compared with rituximab and ofatumumab in vitro and in xenograft models. *Mol Cancer Ther*. 2013;12(10):2031-2042. doi:10.1158/1535-7163.MCT-12-1182 16. Klein C, Lammens A, Schäfer W, et al. Epitope interactions of monoclonal antibodies targeting CD20 and their relationship to functional properties. *mAbs*. 2013;5(1):22-33. doi:10.4161/mabs.22771 17. Mössner E, Brünker P, Moser S, et al. Increasing the efficacy of CD20 antibody therapy through the engineering of a new type II anti-CD20 antibody with enhanced direct and immune effector cell-mediated B-cell cytotoxicity. *Blood*. 2010;115(22):4393-4402. doi:10.1182/blood-2009-06-225979 18. Data on file. Genentech, Inc. 19. Furie RA, Rovin BH, Garg JP, et al. Efficacy and safety of obinutuzumab in active lupus nephritis. *N Engl J Med*. 2025;392(15):1471-1483. doi:10.1056/NEJMoa2410965 20. Furie RA, Aroca G, Cascino MD, et al. B-cell depletion with obinutuzumab for the treatment of proliferative lupus nephritis: a randomised, double-blind, placebo-controlled trial. *Ann Rheum Dis*. 2022;81(1):100-107. doi:10.1136/annrheumdis-2021-220920 21. Costedoat-Chalumeau N, Houssiau FA. Improving medication adherence in patients with lupus nephritis. *Kidney Int*. 2021;99(2):285-287. doi:10.1016/j.kint.2020.10.037 22. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med*. 2005;353(5):487-497. doi:10.1056/NEJMr050100





Actor portrayal.

Discover What GAZYVA + ST* Can Offer Your Patients With Lupus Nephritis¹

- ✓ **Set a higher bar in lupus nephritis with GAZYVA + ST vs placebo + ST¹**
 - 46.4% of patients achieved CRR with GAZYVA + ST vs 33.1% with placebo + ST, with a treatment difference of 13.4% (95% CI, 2.0, 24.8; $P=0.0232$)[†]
 - 42.7% of patients achieved CRR + steroid tapering (≤ 7.5 mg/d) with GAZYVA + ST vs 30.9% with placebo + ST, with a treatment difference of 11.9% (95% CI, 0.6, 23.2; $P=0.0421$)[‡]
- ✓ **The first and only FDA-approved B-cell depleter in lupus nephritis¹**
 - In REGENCY, nearly all patients experienced B-cell depletion by Week 4, sustained through Week 76 with GAZYVA^{§||}
- ✓ **Twice-yearly IV dosing¹**
 - After 4 initial doses, patients receive only 1 dose once every 6 months
- ✓ **Well-demonstrated safety profile^{1,20}**
 - Safety data pooled from 2 clinical studies (REGENCY and NOBILITY)¹
 - The most common ARs ($\geq 5\%$ in the GAZYVA + ST arm) were upper respiratory tract infection, COVID-19, urinary tract infection, bronchitis, pneumonia, infusion-related reactions, and neutropenia¹
 - The most common serious ARs in the GAZYVA + ST arm were COVID-19 (5.5%), pneumonia (4.5%), neutropenia (3.5%), urinary tract infections (2.5%), and infusion-related reactions (0.5%). Two fatal ARs of COVID-19 were reported in the GAZYVA + ST arm¹
- ✓ **EULAR 2025 Guidelines recommend early use of GAZYVA²**
 - For patients with active lupus nephritis, especially those with poor prognostic factors, GAZYVA is a recommended option in combination with standard therapy

GAZYVA has 10+ years of real-world use in oncology.¹

*Standard therapy (ST) included mycophenolate mofetil (MMF) 2 g/d to 2.5 g/d and a tapering course of corticosteroids to a target dose of 5 mg/d.¹
[†]CRR at Week 76: defined as UPCr <0.5 g/g; an eGFR $\geq 85\%$ of baseline, with no occurrence of rescue therapy, treatment failure, death, or early study withdrawal through Week 76.¹
[‡]Defined as maintaining ≤ 7.5 mg/d between Weeks 64 and 76.¹
[§]Defined as B-cell count <10 cells/ μ L. B-cell depletion was observed in 99% of patients by Week 4 and remained below this threshold in 95% of patients at Week 76.¹
^{||}B-cell depletion has not been shown to be directly correlated to clinical response.¹
AR=adverse reaction; CI=confidence interval; CRR=complete renal response; eGFR=estimated glomerular filtration rate; EULAR=European Alliance of Associations for Rheumatology; UPCr=urine protein-to-creatinine ratio.

Indication

GAZYVA® (obinutuzumab) is indicated for the treatment of adult patients with active lupus nephritis (LN) who are receiving standard therapy.

Select Important Safety Information

BOXED WARNINGS: HEPATITIS B VIRUS REACTIVATION AND PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY

- Hepatitis B Virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients receiving CD20-directed cytolytic antibodies, including GAZYVA. Screen all patients for HBV infection before treatment initiation. Monitor HBV-positive patients during and after treatment with GAZYVA. Discontinue GAZYVA and concomitant medications in the event of HBV reactivation
- Progressive Multifocal Leukoencephalopathy (PML), including fatal PML, can occur in patients receiving GAZYVA

Please see additional Important Safety Information throughout and the full [Prescribing Information](#), including BOXED WARNINGS.

Genentech
A Member of the Roche Group

GAZYVA® is a registered trademark of Genentech, Inc.
© 2025 Genentech USA, Inc. All rights reserved.
M-US-00026664(v1.0) 10/25

GAZYVA®
obinutuzumab
injection 1,000mg/40mL



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use GAZYVA safely and effectively. See full prescribing information for GAZYVA.

GAZYVA® (obinutuzumab) injection, for intravenous use

Initial U.S. Approval: 2013

WARNING: HEPATITIS B VIRUS REACTIVATION and PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY

See full prescribing information for complete boxed warning.

- Hepatitis B Virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death. (5.1)
- Progressive Multifocal Leukoencephalopathy (PML) resulting in death. (5.2)

RECENT MAJOR CHANGES

Indication and Usage, Lupus Nephritis (1.3) ----- 10/2025

Dosage and Administration (2.4)----- 10/2025

Warnings and Precautions (5)----- 10/2025

INDICATIONS AND USAGE

GAZYVA is a CD20-directed cytolytic antibody indicated:

- in combination with chlorambucil, for the treatment of patients with previously untreated chronic lymphocytic leukemia (CLL). (1.1, 14)
- in combination with bendamustine followed by GAZYVA monotherapy, for the treatment of patients with follicular lymphoma (FL) who relapsed after, or are refractory to, a rituximab-containing regimen. (1.2, 14)
- in combination with chemotherapy followed by GAZYVA monotherapy in patients achieving at least a partial remission, for the treatment of adult patients with previously untreated stage II bulky, III or IV follicular lymphoma. (1.2, 14)
- for the treatment of adult patients with active lupus nephritis (LN) who are receiving standard therapy (1.3, 14)

DOSAGE AND ADMINISTRATION

- Premedicate for infusion-related reactions and tumor lysis syndrome. (2.1, 5.3, 5.4)
- Administer only as intravenous infusion. Do not administer as an intravenous push or bolus. (2.1)
- The recommended dosage for chronic lymphocytic leukemia is 100 mg on day 1 and 900 mg on day 2 of Cycle 1, 1,000 mg on day 8 and 15 of Cycle 1, and 1,000 mg on day 1 of Cycles 2–6. (2.2)
- The recommended dosage for follicular lymphoma is 1,000 mg on day 1, 8 and 15 of Cycle 1, 1,000 mg on day 1 of Cycles 2–6 or Cycles 2–8, and then 1,000 mg every 2 months for up to 2 years. (2.3)
- The recommended dosage for active lupus nephritis is 1,000 mg at the initial infusion, on Week 2, 24, 26, and every 6 months thereafter. (2.4)

DOSAGE FORMS AND STRENGTHS

Injection: 1,000 mg/40 mL (25 mg/mL) single-dose vial. (3)

CONTRAINDICATIONS

GAZYVA is contraindicated in patients with known hypersensitivity reactions (e.g., anaphylaxis) to obinutuzumab or any of the excipients, including serum sickness with prior obinutuzumab use. (4)

WARNINGS AND PRECAUTIONS

- **Infusion-Related Reactions:** Premedicate patients with glucocorticoid, acetaminophen, and anti-histamine. Monitor patients closely during infusions. Interrupt, reduce rate, or discontinue for infusion-related reactions based on severity. (2.1, 5.3)
- **Hypersensitivity Reactions Including Serum Sickness:** Discontinue GAZYVA permanently. (5.4)
- **Tumor Lysis Syndrome:** In CLL and FL, premedicate with anti-hyperuricemics and adequate hydration, especially for patients with high tumor burden, high circulating lymphocyte count or renal impairment. Correct electrolyte abnormalities, provide supportive care, and monitor renal function and fluid balance. (5.5)
- **Serious, Including Fatal, Infections:** Do not administer GAZYVA to patients with an active infection. Patients with a history of recurring or chronic infections may be at increased risk of infection. (5.6)
- **Neutropenia:** In patients with Grade 3 to 4 neutropenia, monitor laboratory tests until resolution and for infection. Consider dose delays and infection prophylaxis, as appropriate. (5.7)
- **Thrombocytopenia:** Monitor for decreased platelet counts and bleeding. Transfusion may be necessary. (5.8)
- **Disseminated Intravascular Coagulation:** Evaluate cause and monitor for bleeding, thrombosis, and need for supportive care. (5.9)
- **Immunization:** Avoid administration of live virus vaccines during GAZYVA treatment and until B-cell recovery. (5.10)
- **Embryo-Fetal Toxicity:** Can cause fetal harm. Advise females of reproductive potential of the potential risk to a fetus and use effective contraception. (5.11)

ADVERSE REACTIONS

The most common adverse reactions (incidence $\geq 20\%$ and $\geq 2\%$ greater in the GAZYVA treated arm in CLL and NHL, and incidence $\geq 5\%$ in the GAZYVA treated arm in LN) were:

- **Previously untreated CLL:** infusion-related reactions and neutropenia. (6)
- **Relapsed or refractory non-Hodgkin lymphoma (NHL):** infusion-related reactions, fatigue, neutropenia, cough, upper respiratory tract infections, and musculoskeletal pain. (6)
- **Previously untreated NHL:** infusion-related reactions, neutropenia, upper respiratory tract infections, cough, constipation, and diarrhea. (6)
- **Lupus Nephritis:** upper respiratory tract infection, COVID-19, urinary tract infection, bronchitis, pneumonia, infusion-related reactions, and neutropenia. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Genentech at 1-888-835-2555 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- **Lactation:** Advise not to breastfeed. (8.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 10/2025

FULL PRESCRIBING INFORMATION: CONTENTS***WARNING: HEPATITIS B VIRUS REACTIVATION AND PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY****1 INDICATIONS AND USAGE**

- 1.1 Chronic Lymphocytic Leukemia
- 1.2 Follicular Lymphoma
- 1.3 Lupus Nephritis

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosing Information
- 2.2 Recommended Dosage for Chronic Lymphocytic Leukemia
- 2.3 Recommended Dosage for Follicular Lymphoma
- 2.4 Recommended Dosage for Active Lupus Nephritis
- 2.5 Recommended Premedication and Prophylactic Medications
- 2.6 Dosage Modifications for Adverse Reactions
- 2.7 Preparation and Administration

3 DOSAGE FORMS AND STRENGTHS**4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

- 5.1 Hepatitis B Virus Reactivation
- 5.2 Progressive Multifocal Leukoencephalopathy
- 5.3 Infusion-Related Reactions
- 5.4 Hypersensitivity Reactions Including Serum Sickness
- 5.5 Tumor Lysis Syndrome
- 5.6 Infections
- 5.7 Neutropenia
- 5.8 Thrombocytopenia
- 5.9 Disseminated Intravascular Coagulation (DIC)
- 5.10 Immunization
- 5.11 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience in
- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE**11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.6 Immunogenicity

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Chronic Lymphocytic Leukemia
- 14.2 Follicular Lymphoma
- 14.3 Active Lupus Nephritis

16 HOW SUPPLIED/STORAGE AND HANDLING**17 PATIENT COUNSELING INFORMATION**

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: HEPATITIS B VIRUS REACTIVATION and PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY

- **Hepatitis B Virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients receiving CD20-directed cytolytic antibodies, including GAZYVA. Screen all patients for HBV infection before treatment initiation. Monitor HBV-positive patients during and after treatment with GAZYVA. Discontinue GAZYVA and concomitant medications in the event of HBV reactivation [see Warnings and Precautions (5.1)].**
- **Progressive Multifocal Leukoencephalopathy (PML) including fatal PML, can occur in patients receiving GAZYVA [see Warnings and Precautions (5.2)].**

1 INDICATIONS AND USAGE

1.1 Chronic Lymphocytic Leukemia (CLL)

GAZYVA, in combination with chlorambucil, is indicated for the treatment of patients with previously untreated chronic lymphocytic leukemia.

1.2 Follicular Lymphoma (FL)

GAZYVA, in combination with bendamustine followed by GAZYVA monotherapy, is indicated for the treatment of patients with follicular lymphoma who relapsed after, or are refractory to, a rituximab-containing regimen.

GAZYVA, in combination with chemotherapy followed by GAZYVA monotherapy in patients achieving at least a partial remission, is indicated for the treatment of adult patients with previously untreated stage II bulky, III or IV follicular lymphoma.

1.3 Lupus Nephritis (LN)

GAZYVA is indicated for the treatment of adult patients with active lupus nephritis who are receiving standard therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosing Information

- Premedicate before each infusion [see *Dosage and Administration* (2.4)].
- Provide prophylactic hydration and anti-hyperuricemics to patients at high risk of tumor lysis syndrome [see *Dosage and Administration* (2.4) and *Warnings and Precautions* (5.4)].
- Administer only as an intravenous infusion through a dedicated line [see *Dosage and Administration* (2.6)].
- Do not administer as an intravenous push or bolus.
- Monitor blood counts at regular intervals.
- GAZYVA should only be administered by a healthcare professional with appropriate medical support to manage severe infusion-related reactions that can be fatal if they occur [see *Warnings and Precautions* (5.3)].

2.2 Recommended Dosage for Chronic Lymphocytic Leukemia

Each dose of GAZYVA is 1,000 mg administered intravenously with the exception of the first infusions in Cycle 1, which are administered on day 1 (100 mg) and day 2 (900 mg) according to Table 1.

Table 1 Dose of GAZYVA to be Administered During Six 28-Day Treatment Cycles for Patients with CLL

Day of treatment cycle		Dose of GAZYVA	Rate of infusion
Cycle 1 (loading doses)	Day 1	100 mg	Administer at 25 mg/hr over 4 hours. Do not increase the infusion rate.
	Day 2	900 mg	If no infusion-related reaction (IRR) occurred during the previous infusion, administer at 50 mg/hr. The rate of the infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr. If an IRR occurred during the previous infusion, administer at 25 mg/hr. The rate of infusion can be escalated in increments of up to 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.
	Day 8	1,000 mg	If no IRR occurred during the previous infusion and the final infusion rate was 100 mg/hr or faster, infusions can be started at a rate of 100 mg/hr and increased by 100 mg/hr increments every 30 minutes to a maximum of 400 mg/hr.
	Day 15	1,000 mg	
Cycles 2–6	Day 1	1,000 mg	If an infusion-related reaction occurred during the previous infusion, administer at 50 mg/hr. The rate of infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.

If a planned dose of GAZYVA is missed, administer the missed dose as soon as possible and adjust dosing schedule to maintain the time interval between doses. If appropriate, patients who do not complete the Day 1 Cycle 1 dose may proceed to the Day 2 Cycle 1 dose.

2.3 Recommended Dosage for Follicular Lymphoma

Each dose of GAZYVA is 1,000 mg administered intravenously according to Table 2.

For patients with relapsed or refractory FL, administer GAZYVA in combination with bendamustine in six 28-day cycles. Patients who achieve stable disease, complete response, or partial response to the initial 6 cycles should continue on GAZYVA 1,000 mg as monotherapy for up to two years.

For patients with previously untreated FL, administer GAZYVA with one of the following chemotherapy regimens:

- Six 28-day cycles in combination with bendamustine
- Six 21-day cycles in combination with CHOP, followed by 2 additional 21-day cycles of GAZYVA alone
- Eight 21-day cycles in combination with CVP

Patients with previously untreated FL who achieve a complete response or partial response to the initial 6 or 8 cycles should continue on GAZYVA 1,000 mg as monotherapy for up to two years.

GAZYVA should be administered at the standard infusion rate in Cycle 1 (see Table 2). In patients with FL who do not experience a Grade 3 or higher IRR during Cycle 1, GAZYVA may

be administered as a shorter, approximately 90-minute infusion from Cycle 2 onwards (see Table 3) with continued premedication.

Table 2 Dose and Standard Infusion Rate of GAZYVA to be Administered During 6–8 Treatment Cycles, Followed by GAZYVA Monotherapy for Patients with FL

Day of treatment cycle		Dose of GAZYVA	Rate of infusion
Cycle 1 (loading doses)	Day 1	1,000 mg	Administer at 50 mg/hr. The rate of the infusion can be escalated in 50 mg/hr increments every 30 minutes to a maximum of 400 mg/hr.
	Day 8	1,000 mg	If no infusion-related reaction or an infusion-related reaction of Grade 1 occurred during the previous infusion and the final infusion rate was 100 mg/hr or faster, infusions can be started at a rate of 100 mg/hr and increased by 100 mg/hr increments every 30 minutes to a maximum of 400 mg/hr.
	Day 15	1,000 mg	
Cycles 2–6 or 2–8	Day 1	1,000 mg	If an infusion-related reaction of Grade 2 or higher occurred during the previous infusion, administer at 50 mg/hr. The rate of infusion can be escalated in increments of 50 mg/hr every 30 minutes to a maximum rate of 400 mg/hr.
Monotherapy	Every two months for up to two years	1,000 mg	

Table 3 Dose and Infusion Rate of a GAZYVA 90-Minute Infusion for Patients with FL

Day of treatment cycle		Dose of GAZYVA	Rate of infusion
Cycle 1	Days 1, 8, 15	1,000 mg	See Table 2
Cycles 2–6* or 2-8*	Day 1	1,000 mg	If no Grade 3 or higher IRR occurred during Cycle 1: 100 mg/hr for 30 minutes, then 900 mg/hr for approximately 60 minutes. If an IRR of Grade 1-2 with ongoing symptoms or a Grade 3 or higher IRR occurred during the previous approximately 90-minute infusion, administer all subsequent GAZYVA infusions at the standard infusion rate (see Table 2).
Monotherapy*	Every two months for up to two years	1,000 mg	

* Consider an approximately 90-minute infusion in patients with FL who do not experience a Grade 3 or higher infusion-related reaction to GAZYVA in Cycle 1 and subsequent cycles.

If a planned dose of GAZYVA is missed, administer the missed dose as soon as possible. During GAZYVA and chemotherapy treatment, adjust the dosing schedule accordingly to maintain the time interval between chemotherapy cycles. During monotherapy, maintain the original dosing schedule for subsequent doses. Initiate monotherapy approximately two months after the last dose of GAZYVA administered during the induction phase.

2.4 Recommended Dosage for Active Lupus Nephritis

Each recommended dose of GAZYVA is 1,000 mg administered intravenously according to Table 4.

Table 4 Dose and Infusion Rate for Standard Infusion of GAZYVA in Adults with Active Lupus Nephritis

Dose number	Timing of treatment	Dose of GAZYVA	Rate of infusion
1	Initial infusion	1,000 mg	Administer at a rate of 50 mg/hr. The rate of infusion can be escalated in 50 mg/hr increments every 30 minutes to a maximum of 400 mg/hr. For management of IRRs that occur during infusion, see <i>Dosage Modifications for Adverse Reactions (2.6)</i> .
2	Week 2 (two weeks after Dose 1)	1,000 mg	Administer at a rate of 100 mg/hr. The rate of infusion can be escalated at a rate of 100 mg/hr every 30 minutes to a maximum of 400 mg/hr.
3	Week 24	1,000 mg	
4	Week 26 (two weeks after Dose 3)	1,000 mg	
5* and thereafter	Every 6 months	1,000 mg	

*Dose 5 should be administered six months after Dose 4

Patients who do not experience Grade ≥ 3 infusion related reactions during the previous infusion may receive GAZYVA over approximately 90 minutes from Dose 2 onwards (see Table 5), with continued premedication.

Table 5 Dose and Infusion Rate for a 90-Minute Infusion of GAZYVA in Adults with Active Lupus Nephritis

Dose number	Rate of infusion
1	See Table 4
2 and thereafter (if no Grade 3 or higher IRR during previous infusion)	100 mg/hr for 30 minutes, then 900 mg/hr for approximately 60 minutes. If an IRR of Grade 1-2 with ongoing symptoms or a Grade 3 or higher IRR occurs during the previous approximately 90-minute infusion, administer GAZYVA at the standard infusion rate (see Table 4).

If a planned dose of GAZYVA is missed, it should be administered as soon as possible – do not wait until the next planned dose. The schedule of administration should be adjusted to maintain the appropriate interval between doses.

2.5 Recommended Premedication and Prophylactic Medications

Infusion-Related Reactions

Premedication to reduce the risk of IRRs is outlined in Table 6 [see *Warnings and Precautions (5.3)*].

Hypotension may occur during GAZYVA intravenous infusions. Consider withholding antihypertensive treatments for 12 hours prior to and throughout each GAZYVA infusion and for the first hour after administration [see *Warnings and Precautions* (5.3)].

Table 6 Premedication for GAZYVA Infusion to Reduce Infusion-Related Reactions

Indication/Cycle/Day of Treatment	Patients requiring premedication	Premedication	Administration
CLL Cycle 1 Days 1 and 2 FL Cycle 1 Day 1	All patients	Intravenous glucocorticoid: 20 mg dexamethasone or 80 mg methylprednisolone ^{1,2}	Completed at least 1 hour prior to GAZYVA infusion.
		650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.
		anti-histamine (e.g., 50 mg diphenhydramine)	
CLL or FL All subsequent infusions	All patients	650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.
	Patients with an IRR (Grade 1-2) with the previous infusion	650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.
		anti-histamine (e.g., 50 mg diphenhydramine)	
	Patients with a Grade 3 IRR with the previous infusion OR with a lymphocyte count > 25 x 10 ⁹ /L prior to next treatment	Intravenous glucocorticoid: 20 mg dexamethasone or 80 mg methylprednisolone ¹	Completed at least 1 hour prior to GAZYVA infusion.
		650–1,000 mg acetaminophen	At least 30 minutes before GAZYVA infusion.
		anti-histamine (e.g., 50 mg diphenhydramine)	
LN	All patients	Intravenous glucocorticoid (80 mg methylprednisolone)	Completed between 30 and 60 minutes prior to GAZYVA infusion
		650–1,000 mg acetaminophen	Starting from Dose 6, intravenous corticosteroid should only be administered to patients who have experienced an IRR in the prior infusion
		anti-histamine (e.g., 50 mg diphenhydramine)	

Premedication applies to both standard and approximately 90-minute infusions.

¹ Hydrocortisone is not recommended as it has not been effective in reducing the rate of IRRs.

² If a glucocorticoid-containing chemotherapy regimen is administered on the same day as GAZYVA, the glucocorticoid can be administered as an oral medication if given at least 1 hour prior to GAZYVA, in which case additional intravenous glucocorticoid as premedication is not required.

Tumor Lysis Syndrome Prophylaxis

Patients with high tumor burden, high circulating absolute lymphocyte counts (greater than $25 \times 10^9/L$) or renal impairment are considered at risk of tumor lysis syndrome and should receive prophylaxis. Premedicate with anti-hyperuricemics (e.g., allopurinol or rasburicase) and ensure adequate hydration prior to start of GAZYVA therapy. Continue prophylaxis prior to each subsequent GAZYVA infusion, as needed [*see Warnings and Precautions (5.4)*].

Antimicrobial Prophylaxis

Patients with Grade 3 to 4 neutropenia lasting more than one week are strongly recommended to receive antimicrobial prophylaxis until resolution of neutropenia to Grade 1 or 2. Consider antiviral and antifungal prophylaxis for patients with severe and long lasting (> 1 week) neutropenia.

2.6 Dosage Modifications for Adverse Reactions

Infusion-Related Reactions

If a patient experiences an IRR, adjust the infusion as follows [*see Warnings and Precautions (5.3)*]:

- Grade 4 (life-threatening): Stop infusion immediately and permanently discontinue GAZYVA.
- Grade 3 (severe): Interrupt infusion and manage symptoms.
 - For patients who experience Grade 3 IRRs during standard infusion, upon resolution of symptoms, consider restarting GAZYVA infusion at no more than half the previous rate (the rate being used at the time that the IRR reaction occurred), and if patient does not experience any further IRR symptoms, infusion rate escalation may resume at the increments and intervals as appropriate for the treatment cycle dose. Permanently discontinue treatment if patients experience a Grade 3 or higher IRR at rechallenge.
 - For patients with FL who experience Grade 3 IRRs during the approximately 90-minute infusion, upon resolution of symptoms, the infusion can be restarted at no more than half the previous rate (the rate being used at the time that the IRR occurred) and not greater than 400 mg/hr. Administer subsequent infusions at the standard rate. Permanently discontinue treatment if patients experience a Grade 3 or higher IRR at rechallenge.
 - For patients with CLL only, the Day 1 infusion rate may be increased back up to 25 mg/hr after 1 hour but not increased further.
- Grade 1–2 (mild to moderate): Reduce infusion rate or interrupt infusion and manage symptoms. Upon resolution of symptoms, continue or resume GAZYVA infusion, and if patient does not experience any further IRR symptoms, infusion rate escalation may resume at the increments and intervals as appropriate for the treatment cycle dose.
 - For patients with CLL only, the Day 1 infusion rate may be increased back up to 25 mg/hr after 1 hour but not increased further.
 - For patients with LN, reduce infusion rate to half the rate that was used at the time of the reaction and treat symptoms. Upon resolution of symptoms, the infusion should be kept at the reduced rate for an additional 30 minutes. If no further symptoms of IRR, infusion rate escalation may resume at the increments and intervals as appropriate for the treatment dose.

Other Adverse Reactions

Consider treatment interruption if patients experience an infection, Grade 3 or 4 cytopenia, or Grade 2 or higher non-hematologic toxicity.

2.7 Preparation and Administration

Preparation

Prepare the solution for infusion, using aseptic technique, as follows:

- Inspect visually for any particulate matter and discoloration prior to administration.
- Use a sterile needle and syringe to prepare GAZYVA.
- Dilute into a 0.9% Sodium Chloride Injection, USP PVC or non-PVC polyolefin infusion bag.

Chronic Lymphocytic Leukemia

- Preparation of solution for infusion on day 1 (100 mg) and day 2 (900 mg) of Cycle 1:
 - Prepare day 1 (100 mg) and day 2 (900 mg) infusion bags at the same time using one vial (1,000 mg/40 mL) on day 1.
 - Withdraw 40 mL of GAZYVA solution from the vial.
 - Dilute 4 mL (100 mg) of GAZYVA into a 100 mL 0.9% Sodium Chloride Injection, USP infusion bag for immediate administration.
 - Dilute the remaining 36 mL (900 mg) into a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag at the same time for use on day 2 and store in a refrigerator at 2°C to 8°C (36°F to 46°F) for up to 24 hours. After allowing the diluted bag to come to room temperature, use immediately.
 - Clearly label each infusion bag.
- Preparation of solution for infusion on day 8 and 15 of Cycle 1 and day 1 of Cycles 2–6:
 - Withdraw 40 mL of GAZYVA solution from the vial.
 - Dilute 40 mL (1,000 mg) into a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag.

Follicular Lymphoma and Active Lupus Nephritis

- Preparation of solution for infusion:
 - Withdraw 40 mL of GAZYVA solution from the vial.
 - Dilute 40 mL (1,000 mg) into a 250 mL 0.9% Sodium Chloride Injection, USP infusion bag.
- Mix diluted solution by gentle inversion. Do not shake or freeze.
- For microbiological stability, immediately use diluted GAZYVA infusion solution. If not used immediately, store in a refrigerator at 2°C to 8°C (36°F to 46°F) for up to 24 hours prior to use.

The product can be administered at a final concentration of 0.4 mg/mL to 4 mg/mL.

Storage

Use the diluted solution immediately. If not used immediately, store for up to 24 hours at 2°C to 8°C. Discard after 24 hours.

Administration

- Administer as an intravenous infusion only.
- Do not administer as an intravenous push or bolus.
- Do not mix GAZYVA with other drugs.
- No incompatibilities between GAZYVA and polyvinylchloride (PVC) or non-PVC polyolefin bags and administration sets have been observed.

3 DOSAGE FORMS AND STRENGTHS

Injection: 1,000 mg/40 mL (25 mg/mL) clear, colorless to slightly brown solution in a single-dose vial.

4 CONTRAINDICATIONS

GAZYVA is contraindicated in patients with known hypersensitivity reactions (e.g., anaphylaxis) to obinutuzumab or to any of the excipients, or serum sickness with prior obinutuzumab use [*see Warnings and Precautions (5.4)*].

5 WARNINGS AND PRECAUTIONS

5.1 Hepatitis B Virus Reactivation

GAZYVA can cause Hepatitis B virus (HBV) reactivation. Hepatitis B virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients treated with anti-CD20 antibodies such as GAZYVA. HBV reactivation has been reported in patients who are hepatitis B surface antigen (HBsAg) positive and also in patients who are HBsAg negative but are hepatitis B core antibody (anti-HBc) positive. Reactivation has also occurred in patients who appear to have resolved hepatitis B infection (i.e., HBsAg negative, anti-HBc positive, and hepatitis B surface antibody [anti-HBs] positive).

HBV reactivation is defined as an abrupt increase in HBV replication manifesting as a rapid increase in serum HBV DNA level or detection of HBsAg in a person who was previously HBsAg negative and anti-HBc positive. Reactivation of HBV replication is often followed by hepatitis, i.e., increase in transaminase levels and, in severe cases, increase in bilirubin levels, liver failure, and death.

Screen all patients for HBV infection by measuring HBsAg and anti-HBc before initiating treatment with GAZYVA. For patients who show evidence of hepatitis B infection (HBsAg positive [regardless of antibody status] or HBsAg negative but anti-HBc positive), consult healthcare providers with expertise in managing hepatitis B regarding monitoring and consideration for HBV antiviral therapy.

Monitor patients with evidence of current or prior HBV infection for clinical and laboratory signs of hepatitis or HBV reactivation during and for several months following treatment with GAZYVA. HBV reactivation has been reported for other CD20-directed cytolytic antibodies following completion of therapy.

In patients who develop reactivation of HBV while receiving GAZYVA, immediately discontinue GAZYVA and any concomitant chemotherapy and institute appropriate treatment. Resumption of GAZYVA in patients whose HBV reactivation resolves should be discussed with healthcare

providers with expertise in managing hepatitis B. Insufficient data exist regarding the safety of resuming GAZYVA in patients who develop HBV reactivation.

5.2 Progressive Multifocal Leukoencephalopathy

John Cunningham (JC) virus infection resulting in progressive multifocal leukoencephalopathy (PML), which can be fatal, occurred in patients treated with GAZYVA for CLL and NHL. Consider the diagnosis of PML in any patient presenting with new onset or changes to preexisting neurologic manifestations. Evaluation of PML includes, but is not limited to, consultation with a neurologist, brain MRI, and lumbar puncture. Discontinue GAZYVA therapy and consider discontinuation or reduction of any concomitant chemotherapy or immunosuppressive therapy in patients who develop PML.

5.3 Infusion-Related Reactions

GAZYVA can cause severe and life-threatening infusion-related reactions (IRRs). Sixty-five percent of patients with CLL experienced a reaction to the first 1,000 mg of GAZYVA infused. Thirty-seven percent of patients with relapsed or refractory NHL and 60% of patients with previously untreated NHL experienced a reaction on Day 1 of GAZYVA infusion. In patients with CLL and NHL, IRRs have occurred within 24 hours of receiving GAZYVA. IRRs can also occur with subsequent infusions. Symptoms may include hypotension, tachycardia, dyspnea, and respiratory symptoms (e.g., bronchospasm, larynx and throat irritation, wheezing, laryngeal edema). The most frequently reported IRR symptoms in patients with CLL and NHL include nausea, fatigue, chest discomfort, dyspnea, dizziness, vomiting, diarrhea, rash, hypertension, hypotension, flushing, headache, pyrexia, and chills [*see Adverse Reactions (6.1)*].

In patients with LN, IRRs occurred predominantly during infusion of the first 1,000 mg. IRRs were generally mild to moderate and could be managed by the slowing or temporarily halting the infusion [*see Dosage and Administration (2.6)*]. Severe and life-threatening IRRs requiring symptomatic treatment were also reported. The most common IRR signs or symptoms reported in the REGENCY study included headache, nausea and vomiting. In the NOBILITY study, the most common IRR symptoms were pyrexia and tachycardia [*see Adverse Reactions (6.2)*].

Premedicate patients with acetaminophen, anti-histamine, and a glucocorticoid [*see Dosage and Administration (2.4)*]. Closely monitor patients during the entire infusion. Reduce infusion rate, interrupt infusion or permanently discontinue GAZYVA for IRRs based on severity [*see Dosage and Administration (2.5)*]. Institute medical management (e.g., glucocorticoids, epinephrine, bronchodilators, and/or oxygen) for IRRs as needed.

For patients with preexisting cardiac or pulmonary conditions, monitor more frequently throughout the infusion and the post-infusion period since they may be at greater risk of experiencing more severe reactions. Hypotension may occur as part of the IRR to GAZYVA. Consider withholding antihypertensive treatments for 12 hours prior to, during each GAZYVA infusion, and for the first hour after administration until blood pressure is stable. For patients at increased risk of hypertensive crisis, consider the benefits versus the risks of withholding their antihypertensive medication as is suggested here.

5.4 Hypersensitivity Reactions Including Serum Sickness

Hypersensitivity reactions have been reported in patients treated with GAZYVA. Signs of immediate-onset hypersensitivity included dyspnea, bronchospasm, hypotension, urticaria and tachycardia. Late-onset hypersensitivity diagnosed as serum sickness has also been reported, with symptoms that include chest pain, diffuse arthralgia and fever. Hypersensitivity reactions may be difficult to clinically distinguish from IRRs. However, hypersensitivity very rarely occurs with the first infusion and, when observed, often occurs after previous exposure.

If a hypersensitivity reaction is suspected during or after an infusion, stop the infusion and permanently discontinue treatment. GAZYVA is contraindicated in patients with known hypersensitivity reactions to GAZYVA, including serum sickness with prior GAZYVA use [*see Contraindications (4)*].

5.5 Tumor Lysis Syndrome

Tumor lysis syndrome (TLS), including fatal cases, has been reported in patients with CLL and NHL receiving GAZYVA. Patients with high tumor burden, high circulating lymphocyte count ($> 25 \times 10^9/L$) or renal impairment are at greater risk for TLS.

In patients with CLL and NHL at risk for TLS, administer appropriate tumor lysis prophylaxis with anti-hyperuricemics (e.g., allopurinol or rasburicase) and hydration prior to the infusion of GAZYVA [*see Dosage and Administration (2.4)*]. During the initial days of GAZYVA treatment, monitor the laboratory parameters of patients considered at risk for TLS. For treatment of TLS, correct electrolyte abnormalities, monitor renal function and fluid balance, and administer supportive care, including dialysis as indicated. TLS is not identified as a risk in LN.

5.6 Serious, Including Fatal, Infections

Fatal and serious bacterial, fungal, and new or reactivated viral infections can occur during and following GAZYVA therapy.

Lupus Nephritis

In the pooled double-blind periods of REGENCY (Week 76) and NOBILITY (Week 52), the incidence of Grade 3-5 infections was 11% (22/200) in patients treated with GAZYVA and standard therapy compared to 9% (18/193) in patients receiving placebo and standard therapy, corresponding to an exposure-adjusted incidence rate (EAIR) of 8.9 and 7.9 per 100 patient years, respectively. The incidence of fatal infections was 1% (2/200) in patients treated with GAZYVA and 0.5% (1/193) in patients receiving placebo, corresponding to an EAIR of 0.8 and 0.4 per 100 patient years, respectively.

In 40 patients who crossed-over from placebo to GAZYVA and standard therapy at Week 76 in the REGENCY study and patients who continued treatment with GAZYVA and standard therapy, including additional treatment after Week 76, the EAIR of Grade 3-5 infections for the GAZYVA arm was 9.0 per 100 patient-years while the EAIR of fatal infections for the GAZYVA arm was 1.8 per 100 patient-years.

CLL and FL

When GAZYVA is administered with chemotherapy followed by GAZYVA monotherapy as in the GALLIUM study, Grade 3 to 5 infections have been reported in up to 8% of patients during combination therapy, up to 13% of patients during monotherapy, and up to 8% of patients after treatment [*see Adverse Reactions (6.1)*].

In GALLIUM, more Grade 3 to 5 infections were reported in the recipients of GAZYVA and bendamustine (117/410 patients, 29%) as compared to GAZYVA plus CHOP or CVP (43/281 patients, 15%). More fatal infections were reported in patients treated with GAZYVA and bendamustine (3%), as compared to GAZYVA plus CHOP or CVP ($< 1\%$), including during the monotherapy phase and after completion of treatment.

Do not administer GAZYVA to patients with an active infection. Patients with a history of recurring or chronic infections may be at increased risk of infection. In patients who develop a serious infection while receiving GAZYVA, immediately discontinue GAZYVA and institute appropriate treatment. Consider the risk and benefit of resuming treatment with GAZYVA following resolution of serious infections.

5.7 Neutropenia

Severe and life-threatening neutropenia, including febrile neutropenia, has been reported during treatment with GAZYVA. Monitor patients with Grade 3 to 4 neutropenia frequently with regular laboratory tests until resolution. Anticipate, evaluate, and treat any symptoms or signs of developing infection. Consider dose delays for Grade 3 or 4 neutropenia. Consider administration of granulocyte colony-stimulating factors (GCSF) in patients with Grade 3 or 4 neutropenia.

Neutropenia can also be of late onset (occurring more than 28 days after completion of treatment) and/or prolonged (lasting longer than 28 days). Patients with severe and long lasting (> 1 week) neutropenia are strongly recommended to receive antimicrobial prophylaxis until resolution of neutropenia to Grade 1 or 2. Consider antiviral and antifungal prophylaxis.

5.8 Thrombocytopenia

Severe and life-threatening thrombocytopenia has been reported during treatment with GAZYVA in combination with chemotherapy. Fatal hemorrhagic events have been reported in patients with NHL and CLL treated with GAZYVA in combination with chemotherapy, including during Cycle 1.

Monitor patients frequently for thrombocytopenia and hemorrhagic events, especially during the first cycle and if clinically indicated, evaluate laboratory coagulation parameters [*see Warnings and Precautions (5.9)*]. In patients with Grade 3 or 4 thrombocytopenia, monitor platelet counts more frequently until resolution and consider dose delays of GAZYVA and chemotherapy or dose reductions of chemotherapy. Transfusion of blood products (i.e., platelet transfusion) may be necessary. Consider withholding concomitant medications that may increase bleeding risk (platelet inhibitors, anticoagulants), especially during the first cycle.

5.9 Disseminated Intravascular Coagulation (DIC)

Fatal and severe DIC has been reported in patients receiving GAZYVA for treatment of CLL and NHL. The majority of DIC cases have involved changes in platelets and laboratory coagulation parameters following the first infusion, with spontaneous resolution usually occurring by Day 8. In some cases, DIC was associated with IRRs, TLS, or both [*see Adverse Reactions (6.1)*].

In patients with suspected DIC, evaluate potential causes, and monitor coagulation parameters, platelet counts, and for signs and symptoms of bleeding or thrombosis. Manage according to standard guidelines for DIC. Supportive care, including transfusion of blood products and other medical management, may be necessary.

5.10 Immunization

The safety and efficacy of immunization with live or attenuated viral vaccines during or following GAZYVA therapy have not been studied. Immunization with live virus vaccines is not recommended during treatment with GAZYVA and until B-cell recovery.

5.11 Embryo-Fetal Toxicity

Based on its mechanism of action and findings in animals, GAZYVA can cause B-cell depletion in infants exposed to obinutuzumab in-utero. Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception while taking GAZYVA and for 6 months after the last dose [*see Use in Specific Populations (8.1, 8.3)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Hepatitis B virus reactivation [see *Warnings and Precautions* (5.1)]
- Progressive multifocal leukoencephalopathy [see *Warnings and Precautions* (5.2)]
- Infusion-related reactions [see *Warnings and Precautions* (5.3)]
- Hypersensitivity reactions including serum sickness [see *Warnings and Precautions* (5.4)]
- Tumor lysis syndrome [see *Warnings and Precautions* (5.5)]
- Infections [see *Warnings and Precautions* (5.6)]
- Neutropenia [see *Warnings and Precautions* (5.7)]
- Thrombocytopenia [see *Warnings and Precautions* (5.8)]
- Disseminated intravascular coagulation [see *Warnings and Precautions* (5.9)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Chronic Lymphocytic Leukemia

The data below are based on a safety population of 773 previously untreated patients with CLL in the CLL11 study. Patients were treated with chlorambucil alone, GAZYVA in combination with chlorambucil, or rituximab product in combination with chlorambucil. The Stage 1 analysis compared GAZYVA in combination with chlorambucil vs. chlorambucil alone and Stage 2 compared GAZYVA in combination with chlorambucil vs. rituximab product in combination with chlorambucil. Adverse reactions rates and laboratory abnormalities from the Stage 2 phase are presented below and are consistent with the rates in Stage 1. In addition to the adverse reactions observed in Stage 2, in Stage 1, back pain (5% vs. 2%), anemia (12% vs. 10%) and cough (10% vs. 7%) were observed at a higher incidence in the GAZYVA treated patients. The incidence of Grade 3 to 4 back pain (< 1% vs. 0%), cough (0% vs. < 1%) and anemia (5% vs. 4%) was similar in both treatment arms. With regard to laboratory abnormalities, in Stage 1 hyperkalemia (33% vs. 18%), creatinine increased (30% vs. 20%) and alkaline phosphatase increased (18% vs. 11%) were observed at a higher incidence in patients treated with GAZYVA with similar incidences of Grade 3 to 4 abnormalities between the two arms.

Patients received three 1,000 mg doses of GAZYVA on the first cycle and a single dose of 1,000 mg once every 28 days for 5 additional cycles in combination with chlorambucil (6 cycles of 28 days each in total). In the last 140 patients enrolled, the first dose of GAZYVA was split between day 1 (100 mg) and day 2 (900 mg) [see *Dosage and Administration* (2.2)]. In total, 81% of patients received all 6 cycles (of 28 days each) of GAZYVA-based therapy.

Adverse reactions in $\geq 10\%$ of patients in the GAZYVA containing arm were infusion-related reactions, neutropenia, thrombocytopenia, and diarrhea. The most common Grade 3 to 4 adverse reactions (incidence $\geq 10\%$) in the GAZYVA containing arm were neutropenia, infusion-related reactions, and thrombocytopenia.

Table 7 Adverse Reactions (Incidence $\geq 5\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with CLL (Stage 2)

Body System Adverse Reactions	GAZYVA + Chlorambucil n = 336		Rituximab product + Chlorambucil n = 321	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Injury, Poisoning and Procedural Complications				
Infusion-Related Reaction	66	20	38	4
Blood and Lymphatic System Disorders^a				
Neutropenia	38	33	32	28
Thrombocytopenia	14	10	7	3
Gastrointestinal Disorders				
Diarrhea	10	2	8	< 1
Constipation	8	0	5	0
General Disorders and Administration Site Conditions				
Pyrexia	9	< 1	7	< 1
Infections and Infestations				
Nasopharyngitis	6	< 1	3	0
Urinary Tract Infection	5	1	2	< 1

^a Adverse reactions reported under “Blood and lymphatic system disorders” reflect those reported by investigator as clinically significant.

Table 8 Post-Baseline Laboratory Abnormalities (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with CLL (Stage 2)

Laboratory Abnormalities	GAZYVA + Chlorambucil n = 336		Rituximab product + Chlorambucil n = 321	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Hematology				
Leukopenia	84	35	62	16
Lymphopenia	80	39	50	16
Neutropenia	76	46	69	41
Thrombocytopenia	48	13	40	8
Anemia	39	10	37	10
Chemistry				
Hypocalcemia	37	3	32	< 1
ALT increased	28	2	21	1
AST increased	27	2	21	< 1
Hyponatremia	26	7	18	2
Hypoalbuminemia	23	< 1	16	< 1
Hypokalemia	14	1	10	< 1

Non-Hodgkin Lymphoma

GADOLIN Study

The GADOLIN study evaluated safety in 407 patients with relapsed or refractory NHL, including FL (81%), small lymphocytic lymphoma and marginal zone lymphoma (a disease for which GAZYVA is not indicated), who did not respond to or progressed within 6 months of treatment with rituximab product or a rituximab product-containing regimen. In the population of patients with FL, the profile of adverse reactions was consistent with the overall NHL population. Patients received either GAZYVA in combination with bendamustine (204 patients), followed by GAZYVA monotherapy in patients that had not progressed, or bendamustine alone (203 patients).

Patients randomized to the GAZYVA + bendamustine arm received three weekly 1,000 mg doses of GAZYVA in the first cycle and a single dose of 1,000 mg once every 28 days for 5 additional cycles, in combination with bendamustine 90 mg/m² intravenously on Days 1 and 2 in all 6 cycles. Patients who did not progress on the combination received a single 1,000 mg dose of GAZYVA

monotherapy every two months until progression or for a maximum of two years. The control arm received bendamustine 120 mg/m² on Days 1 and 2 of each cycle for 6 cycles, with a cycle length of 28 days. In the GAZYVA arm, 78% of patients received 6 cycles of bendamustine and 82% received their full 6 cycles of GAZYVA; 72 (46%) of the 158 patients who began GAZYVA monotherapy received all planned doses. In the control arm, 72% of patients received 6 cycles of bendamustine.

Serious adverse reactions occurred in 45% of the GAZYVA arm and 37% of the bendamustine-only arm. Fatal adverse reactions within 90 days of treatment occurred in 3.4% and 2.5%, respectively. During treatment and follow-up combined, fatal adverse reactions occurred in 10% of GAZYVA recipients and in 7.4% of recipients of bendamustine alone, with infection and second primary malignancies being the leading causes.

Dose modification due to adverse reactions occurred in 50% of the GAZYVA arm and 42% of the control arm, and discontinuation of any study drug due to adverse reactions occurred in 20% and 17%, respectively.

Table 9 presents selected adverse reactions in GADOLIN. The most common adverse reactions (incidence $\geq$ 20%) in GAZYVA recipients included infusion-related reactions, fatigue, neutropenia, cough, upper respiratory tract infections, and musculoskeletal pain.

Table 9 Adverse Reactions (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with Relapsed or Refractory NHL (GADOLIN)

Body System Adverse Reactions ^{a, b}	GAZYVA + Bendamustine followed by GAZYVA monotherapy n = 204		Bendamustine n = 203	
	All Grades %	Grades 3 to 5 %	All Grades %	Grades 3 to 5 %
Procedural Complications				
Infusion-Related Reaction ^c	67	11	63	5
General Disorders				
Fatigue	40	3	36	3
Pyrexia	19	1	15	1
Blood and Lymphatic System Disorders				
Neutropenia	37	35 ^d	29	27
Infections and Infestations				
Upper Respiratory Tract Infection	36	3	23	1
Respiratory Tract Infection, Unspecified	14	1	8	0
Urinary Tract Infection	13	3	7	0
Respiratory, Thoracic and Mediastinal Disorders				
Cough	31	<1	21	0
Musculoskeletal and Connective Tissue Disorders				
Musculoskeletal Pain	28	1	20	0
Arthralgia	12	<1	5	0
Skin and Subcutaneous Tissue Disorders				
Rash	17	<1	14	<1
Pruritus	11	0	6	0

^a Includes adverse reactions reported throughout study treatment and follow-up.

^b Includes grouped terms.

^c Except where noted, individual events that meet the definition of “infusion-related reaction” are excluded from Table 9 above, as they are included in the grouped term “Infusion-Related Reaction”.

^d Includes 1 fatal event.

Infusion-related reactions are defined as any related adverse reaction that occurred during or within 24 hours of infusion.

Fatigue includes fatigue, lethargy, asthenia.

Pyrexia includes pyrexia, hyperthermia, body temperature increased.

Cough includes cough, productive cough, upper-airway cough syndrome.

Neutropenia includes neutropenia, agranulocytosis, granulocytopenia, neutrophil count decreased.

Upper respiratory tract infection includes upper respiratory tract congestion, upper respiratory tract inflammation, upper respiratory fungal infection, rhinovirus infection, and all terms containing: upper respiratory tract infection, laryngitis, nasopharyngitis, pharyngitis, rhinitis, tonsillitis, and sinusitis with the exception of sinobronchitis.

Respiratory tract infection unspecified includes respiratory tract infection, respiratory tract infection viral, influenza, influenza-like illness, sinobronchitis, respiratory syncytial virus infection.

Urinary tract infection includes all terms containing: urinary tract infection, cystitis, pyelonephritis.

Musculoskeletal pain includes non-cardiac chest pain, bone pain, spinal pain, myalgia, back pain, neck pain, musculoskeletal discomfort, pain in extremity, and all terms containing “musculoskeletal pain”.

Rash includes drug eruption, skin reaction, all terms containing “rash”, urticaria, and selected terms containing “dermatitis”.

Pruritus includes pruritus, pruritus generalized.

Other clinically relevant adverse reactions (incidence < 10% and $\geq 2\%$ greater in the GAZYVA arm) included:

- *Blood and lymphatic system disorders*: febrile neutropenia (6%)
- *Infection*: sepsis (7%)

During GAZYVA monotherapy (158 patients), adverse reactions in $\geq 10\%$ of patients included upper and lower respiratory tract infections, cough, neutropenia, musculoskeletal pain, fatigue, diarrhea, rash, and urinary tract infection.

Table 10 presents selected new or worsening laboratory abnormalities in the GADOLIN trial.

Table 10 New or Worsening Laboratory Abnormalities (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm^a) in Patients with Relapsed or Refractory NHL (GADOLIN)

Laboratory Abnormalities	GAZYVA + Bendamustine followed by GAZYVA monotherapy n = 204		Bendamustine n = 203	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Hematology				
Lymphopenia	97	92	96	84
Leukopenia	84	47	87	34
Neutropenia	76	53	75	42
Chemistry				
Hypophosphatemia	41	8	38	7
Hypocalcemia	39	3	24	1
ALT/SGPT increased	36	2	31	3
Alkaline phosphatase increased	27	0	23	0
Hyperbilirubinemia	21	2	17	2
Hyperkalemia	20	3	18	0

^a Two percent difference in either any-grade or Grade 3 to 4 laboratory abnormalities.

In the GAZYVA monotherapy phase, new or worsening grade 3 or 4 abnormalities included neutropenia in 25% of patients (Grade 4, 10%) and lymphopenia in 23% (Grade 4, 5%).

GALLIUM Study

A randomized, open-label multicenter trial (GALLIUM) evaluated the safety of GAZYVA as compared to rituximab product in 1385 patients with previously untreated follicular lymphoma (86%) or marginal zone lymphoma (14%). Patients received chemotherapy (bendamustine, CHOP, or CVP) combined with either GAZYVA (691 patients) or rituximab product (694 patients), followed in responding patients by GAZYVA or rituximab product monotherapy every two months until disease progression or for a maximum of two years. The study excluded patients having an absolute neutrophil count (ANC) $< 1500 / \mu\text{L}$, platelets $< 75,000 / \mu\text{L}$, CLcr $< 40 \text{ mL/min}$ and, unless attributable to lymphoma, hepatic transaminases $> 2.5 \times$ upper limit of normal.

The median age was 60 (range: 23-88), 47% were male, 82% were white, and 97% had an ECOG performance status of 0 or 1. The chemotherapy was bendamustine in 59%, CHOP in 31% and CVP in 10% of patients. Following combination therapy, 624 patients (90%) in the GAZYVA arm and 612 patients (88%) in the rituximab product arm received monotherapy.

Serious adverse reactions occurred in 50% of patients on the GAZYVA arm and 43% of patients on the rituximab product arm. Fatal adverse reactions were reported during treatment in 3% in the GAZYVA arm and 2% in the rituximab product arm, most often from infections in the GAZYVA arm. During treatment and follow-up combined, fatal adverse reactions were reported in 5% of the GAZYVA arm and 4% of the rituximab product arm, with infections and second malignancies being leading causes. In the GAZYVA arm, fatal infections occurred in 2% of patients compared to $< 1\%$ in the rituximab product arm.

During combination therapy, 93% of patients received all treatment cycles in the GAZYVA arm, and 92% received all treatment cycles in the rituximab product arm. Of the responding patients who began monotherapy with GAZYVA or rituximab product, 76% and 73%, respectively, completed the full course. Dose modification due to adverse reactions occurred in 74% of the GAZYVA arm and 63% of the rituximab product arm throughout study treatment, and discontinuation of any study drug due to adverse reactions occurred in 18% and 15%, respectively.

Throughout treatment and follow-up, the most common adverse reactions (incidence $\geq 20\%$) observed at least 2% more in the GAZYVA arm included infusion-related reactions, neutropenia, upper respiratory tract infections, constipation and diarrhea (Table 11). Neutropenia, infusion-related

reactions, febrile neutropenia and thrombocytopenia were the most common Grade 3 to 5 adverse reactions (incidence $\geq 5\%$) observed more frequently in the GAZYVA arm.

Table 11 Adverse Reactions (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with Previously Untreated NHL (GALLIUM)

Body System Adverse Reactions ^{a, b}	GAZYVA + chemotherapy followed by GAZYVA monotherapy n = 691		Rituximab product + chemotherapy followed by rituximab product monotherapy n = 694	
	All Grades %	Grades 3 to 5 %	All Grades %	Grades 3 to 5 %
Injury, Poisoning and Procedural Complications				
Infusion-Related Reaction ^c	72	12	60	8
Blood and Lymphatic System Disorders				
Neutropenia ^d	53	49	47	41
Thrombocytopenia ^d	14	7	8	3
Infections and Infestations				
Upper Respiratory Tract Infection	50	3	43	1
Herpesvirus Infection	18	3	14	1
Pneumonia	14	7	12	6
Respiratory, Thoracic and Mediastinal Disorders				
Cough	35	< 1	28	< 1
Gastrointestinal Disorders				
Constipation	32	< 1	29	< 1
Diarrhea	30	3	26	2
Nervous System Disorders				
Headache	18	< 1	15	< 1
Musculoskeletal and Connective Tissue Disorders				
Arthralgia	16	0	14	< 1
Psychiatric Disorders				
Insomnia	15	< 1	12	< 1
Metabolism and Nutrition Disorders				
Decreased Appetite	14	< 1	12	< 1
Skin and Subcutaneous Tissue Disorders				
Pruritus	11	< 1	9	0

^a Includes adverse reactions reported throughout study treatment and follow-up.

^b Includes grouped terms.

^c Except where noted, individual events that meet the definition of “infusion-related reaction” are excluded from Table 11 above, as they are already included in the grouped term “Infusion-Related Reaction”. The most common individual terms within the grouped term “Infusion-Related Reaction” in decreasing order of frequency are nausea, chills, pyrexia and vomiting.

^d Includes adverse reactions reported as infusion-related reactions.

Infusion-related reactions are defined as any related adverse reaction that occurred during or within 24 hours of infusion.

Neutropenia includes neutropenia, agranulocytosis, granulocytopenia, and neutrophil count decreased.

Febrile neutropenia includes febrile neutropenia, neutropenic infection, neutropenic sepsis, and febrile bone marrow aplasia.

Thrombocytopenia includes thrombocytopenia and platelet count decreased.

Upper respiratory tract infection includes upper respiratory tract congestion, upper respiratory tract inflammation, upper respiratory tract infection, rhinovirus infection, and all terms containing: laryngitis, nasopharyngitis, pharyngitis, rhinitis, tonsillitis, and sinusitis with the exception of sinobronchitis.

Herpesvirus infection includes all terms containing “herpes” or “varicella.”

Pneumonia includes all terms containing “pneumonia,” bacterial, pneumonia haemophilus, pneumonia pneumococcal, pneumonia fungal, pneumocystis jirovecii infection, lung infection, and lung infiltration.

Diarrhea includes diarrhea, defecation urgency, frequent bowel movement, and all terms containing “gastroenteritis”.

Headache includes all terms containing “headache” and migraine.

Insomnia includes all terms containing “insomnia” and sleep disorder.

Pruritus includes pruritus and pruritus generalized.

During the monotherapy period, the common adverse reactions (incidence $\geq 10\%$) observed at least 2% more with GAZYVA were upper respiratory tract infection (40%), cough (23%), musculoskeletal pain (20%), neutropenia (19%), and herpesvirus infection (13%).

Table 12 summarizes treatment-emergent laboratory abnormalities during treatment and follow-up. The Grade 3 to 4 abnormalities reported at least 2% more in the GAZYVA arm were lymphopenia, leukopenia, neutropenia, thrombocytopenia, and hyperuricemia. Patients in the GAZYVA arm, as compared to the rituximab product arm, had higher incidences of Grade 4 neutropenia (38% vs. 30%, respectively), Grade 4 lymphopenia (33% vs. 22%), and Grade 4 leukopenia (17% vs. 12%).

Table 12 New or Worsening Laboratory Abnormalities (Incidence $\geq 10\%$ and $\geq 2\%$ Greater in the GAZYVA Arm) in Patients with Previously Untreated NHL (GALLIUM)

Laboratory Abnormalities ^a	GAZYVA+ chemotherapy followed by GAZYVA monotherapy n = 691		Rituximab product + chemotherapy followed by rituximab product monotherapy n = 694	
	All Grades %	Grades 3 to 4 %	All Grades %	Grades 3 to 4 %
Hematology				
Lymphopenia	97	83	95	67
Leukopenia	92	49	89	39
Neutropenia	84	59	76	50
Thrombocytopenia	68	11	50	4
Chemistry				
ALT increased	50	3	43	2
AST increased	44	1	41	1
Hypophosphatemia	36	5	33	5
Hypoalbuminemia	33	1	25	1
Hypoproteinemia	32	0	30	0
Hypocalcemia	32	1	26	1
Hyperuricemia	28	28	22	22
Hyponatremia	26	4	20	3
Hyperkalemia	23	1	17	1
Hypernatremia	16	< 1	13	0

^a Includes lab abnormalities, reported throughout treatment and follow-up, that were new or worsening, or worsening from baseline unknown.

In the monotherapy phase, new-onset Grade 3 or 4 neutropenia was reported in 21% of patients in the GAZYVA arm (Grade 4, 10%) and 17% of patients in the rituximab product arm (Grade 4, 9%).

GAZELLE Study

GAZELLE (NCT03817853) is a single-arm study designed to characterize the safety of GAZYVA administered as a shortened-duration infusion (approximately 90 minutes) in patients with previously untreated FL. All patients received GAZYVA at the standard infusion rate with premedication during Cycle 1. If no Grade 3 or higher IRR occurred with any infusion during Cycle 1, GAZYVA was administered over approximately 90 minutes in Cycle 2 and subsequent cycles. The primary safety measure was the proportion of patients who experienced Grade 3 or higher IRRs with the 90-minute infusion at Cycle 2. GAZYVA was administered in combination with CHOP, CVP or bendamustine for 6 to 8 cycles (induction), followed by monotherapy for up to 2 years.

Of the 113 patients treated with GAZYVA, 99 (88%) received the 90-minute infusion starting in Cycle 2. In total, 97% of patients who received GAZYVA at either a standard or shorter infusion duration received premedication in Cycle 2. IRRs were observed in 63% of patients throughout induction (including IRRs observed after standard-duration infusion). In cycle 1, 58% of patients developed IRRs with the standard infusion rate (Grade ≥ 3 IRR, 5%). Of the patients who received the 90-minute infusion, 10% had IRRs of any grade in Cycle 2, with 8% and 2% of patients having a

Grade 1 IRR or Grade 2 IRR, respectively. Following Cycle 2, one (1%) patient experienced a Grade 3 IRR, which occurred after the 90-minute infusion at Cycle 5.

Lupus Nephritis

The data below reflects exposure to GAZYVA administered intravenously in patients with ISN/RPS 2003 Class III or IV with or without concomitant Class V lupus nephritis in the REGENCY and NOBILITY studies, up to week 76.

REGENCY (NCT04221477) is a Phase III study which included 136 patients treated with GAZYVA plus standard therapy consisting of mycophenolate mofetil (MMF) and corticosteroids [*see Clinical Studies (14.3)*].

NOBILITY (NCT02550652) is a Phase II study which included 64 patients treated with GAZYVA plus standard therapy consisting of MMF/mycophenolic acid (MPA) and corticosteroids.

The most common adverse reactions (incidence $\geq 5\%$ in the GAZYVA arm) were upper respiratory tract infection, COVID-19, urinary tract infection, bronchitis, pneumonia, infusion related reactions and neutropenia.

The most common serious adverse reactions in the GAZYVA arm were: COVID-19 (11 patients [5.5%]), pneumonia (9 patients [4.5%]), neutropenia (7 patients [3.5%]), urinary tract infections (5 patients [2.5%]) and infusion-related reactions (1 patient [0.5%]). No serious adverse reactions were reported for bronchitis, herpes simplex and upper respiratory tract infections. Two fatal adverse reactions of COVID-19 were reported in the GAZYVA arm.

Adverse reaction rates are presented in Table 13.

Table 13 Adverse Reactions in Adults with Active Lupus Nephritis Treated with GAZYVA Versus Placebo ($\geq 5\%$ Greater in the GAZYVA Arm) in REGENCY and NOBILITY Studies

Body System Adverse Reactions	GAZYVA + standard therapy n = 200		Placebo + standard therapy n = 193	
	All Grades (%)	Grades 3- 5(%)	All Grades (%)	Grades 3-5 (%)
<u>Infections and Infestations</u>				
Upper Respiratory Tract Infection	29	0	24	0
COVID-19	23	5 ^a	16	0.5 ^a
Urinary Tract Infection	21	3	18	4
Bronchitis	14	0	8	0.5
Pneumonia	10	2	6	2
Herpes Simplex	3	0	0	0
<u>Injury, Poisoning and Procedural Complications</u>				
Infusion Related Reaction	14	2	10	0.5
<u>Blood and Lymphatic System Disorders</u>				
Neutropenia	14	7	6	0.5

^a Includes 2 fatal events in the GAZYVA arm and 1 fatal event in the placebo arm

Specific Adverse Reactions

Infusion-Related Reactions

Chronic Lymphocytic Leukemia

The incidence of IRRs in the CLL11 study was 65% with the first infusion of GAZYVA. The incidence of Grade 3 or 4 IRRs was 20% with 7% of patients discontinuing therapy. The incidence of reactions with subsequent infusions was 3% with the second 1,000 mg and < 1% thereafter. No Grade 3 or 4 IRRs were reported beyond the first 1,000 mg infused.

Of the first 53 patients receiving GAZYVA in CLL11, 47 (89%) experienced an IRR. After this experience, study protocol modifications were made to require pre-medication with a corticosteroid, antihistamine, and acetaminophen. The first dose was also divided into two infusions (100 mg on day 1 and 900 mg on day 2). For the 140 patients for whom these mitigation measures were implemented, 74 patients (53%) experienced a reaction with the first 1,000 mg (64 patients on day 1, 3 patients on day 2, and 7 patients on both days) and < 3% thereafter [see *Dosage and Administration* (2.2)].

Non-Hodgkin Lymphoma

Overall, 67% of patients in the GADOLIN study experienced an IRR (all grades) during treatment with GAZYVA in combination with bendamustine. The incidence of Grade 3 to 4 IRRs in GADOLIN was 11%. In Cycle 1, the incidence of IRRs (all grades) was 53% in patients receiving GAZYVA in combination with bendamustine of which 34 (9%) were Grade 3 to 4 in severity. In patients receiving GAZYVA in combination with bendamustine, the incidence of IRRs was highest on Day 1 (37%), and gradually decreased on Days 2, 8 and 15 (23%, 6% and 4%, respectively).

During Cycle 2, the incidence of IRRs was 24% in patients receiving GAZYVA in combination with bendamustine and decreased with subsequent cycles.

During GAZYVA monotherapy in GADOLIN, IRRs (all grades) were observed in 8% of patients. One Grade 3 and no Grade 4 IRRs were reported during GAZYVA monotherapy.

Overall, 2% of patients in GADOLIN experienced an IRR leading to discontinuation of GAZYVA.

In GALLIUM, 72% of patients in the GAZYVA treated arm experienced an IRR (all grades). The incidence of Grade 3 to 4 IRRs for these patients was 12%. In Cycle 1, the incidence of IRRs (all grades) was 62% in the GAZYVA treated arm with Grade 3 to 4 IRRs reported in 10%. The incidence of IRRs (all grades) was highest on Day 1 (60%) and decreased on Days 8 and 15 (9% and 6%, respectively).

During Cycle 2, the incidence of IRRs (all grades) in the GAZYVA treated arm was 13% and decreased with subsequent cycles.

During GAZYVA monotherapy treatment in GALLIUM, IRRs (all grades) were observed in 9% of patients.

Overall, 1% of patients in GALLIUM experienced an IRR leading to discontinuation of GAZYVA.

In GAZELLE, 10% of patients with FL experienced IRRs of any grade at Cycle 2 when GAZYVA was administered over approximately 90 minutes.

Lupus Nephritis

In the pooled REGENCY and NOBILITY studies, infusion-related reactions (IRRs) were reported in 14% of patients in the GAZYVA arm versus 10% of patients in the placebo arm. IRRs in both arms were predominantly Grade 1-2 and occurred during or after the first infusion. Grade 3-4 IRRs were reported in 1.5% of patients in the GAZYVA arm vs 0.5% of patients in the placebo arm. All Grade 3-4 events occurred during or after either the first or second infusion.

The incidence of IRRs in the GAZYVA arm decreased from 11% during the first infusion to 3% during the second infusion, decreasing further with subsequent infusions to 0.5% during the sixth infusion. The severity of IRRs in the GAZYVA arm also decreased with subsequent infusions with 1% patients reporting Grade 3-4 IRRs during the first infusion and 0.5% patients reporting Grade 3-4 IRRs during the second infusion. In subsequent infusions all IRRs were Grade 1-2 in severity. No Grade 5 IRRs were reported [see *Warnings and Precautions* (5.3)].

In the REGENCY study, most common IRR signs or symptoms included headache, nausea and vomiting. In the NOBILITY study, the most common IRR symptoms were pyrexia and tachycardia.

Neutropenia

Chronic Lymphocytic Leukemia

The incidence of neutropenia reported as an adverse reaction in CLL11 was 38% in the GAZYVA treated arm and 32% in the rituximab product treated arm, with the incidence of serious adverse reactions being 1% and < 1%, respectively (Table 7). Cases of late-onset neutropenia (occurring 28 days after completion of treatment or later) were 16% in the GAZYVA treated arm and 12% in the rituximab product treated arm.

Non-Hodgkin Lymphoma

The incidence of neutropenia in GADOLIN was higher in the GAZYVA plus bendamustine arm (37%) compared to the arm treated with bendamustine alone (30%). Cases of prolonged neutropenia (3%) and late onset neutropenia (8%) were also reported in the GAZYVA plus bendamustine arm. The incidence of neutropenia was higher during treatment with GAZYVA in combination with bendamustine (30%) compared to the GAZYVA monotherapy treatment phase (13%).

The incidence of neutropenia in GALLIUM was higher in the GAZYVA treated arm (53%) compared to the rituximab product treated arm (47%). Cases of prolonged neutropenia (1%) and late onset neutropenia (4%) were also reported in the GAZYVA treated arm. The incidence of neutropenia was higher during treatment with GAZYVA in combination with chemotherapy (45%) compared to the GAZYVA monotherapy treatment phase (20%).

Lupus Nephritis

In the pooled REGENCY and NOBILITY studies, neutropenia and related adverse reactions (i.e., leukopenia, lymphopenia, lymphocyte count decreased, febrile neutropenia, and neutrophil count decreased) were reported in 14% of patients in the GAZYVA arm versus 6% of patients in the placebo arm. Grade 3-4 neutropenia was reported in 7% of patients treated with GAZYVA versus 0.5% of patients in the placebo arm. Neutropenia and related adverse reactions resolved/improved spontaneously or with use of granulocyte colony-stimulating factors in 96% of patients [see *Warnings and Precautions (5.7)*].

Infection

Chronic Lymphocytic Leukemia

The incidence of infections was similar between GAZYVA and rituximab product treated arms. Thirty-eight percent of patients in the GAZYVA treated arm and 37% in the rituximab product treated arm experienced an infection, with Grade 3 to 4 rates being 11% and 13%, respectively. Fatal events were reported in 1% of patients in both arms.

Non-Hodgkin Lymphoma

The incidence of infection in GADOLIN was 68% in the GAZYVA plus bendamustine arm and 59% in the bendamustine arm, with Grade 3 to 4 events reported in 20% and 16%, respectively. Fatal events were reported in 3% of patients in the GAZYVA plus bendamustine arm and 3% in the bendamustine arm.

The incidence of infections in GALLIUM was 82% in the GAZYVA treated arm and 73% in the rituximab product treated arm, with Grade 3 to 4 events reported in 21% and 17%, respectively. In the GAZYVA arm, fatal infections occurred in 2% of patients compared to <1% in the rituximab product arm.

The incidence of Grade 3 to 4 infections in the GAZYVA and rituximab product treated arms was lower in patients receiving GCSF prophylaxis (14%; 16%) compared with patients not receiving GCSF prophylaxis (24%; 18%). The incidence of fatal infections in patients receiving GCSF prophylaxis in the GAZYVA and rituximab product treated arms was 2% and 0%, respectively, and was 2% and < 1% in patients not receiving GCSF prophylaxis.

Lupus Nephritis

In the pooled REGENCY and NOBILITY studies, infections were reported in 72% of patients in the GAZYVA arm versus 62% of patients in the placebo arm. The most frequently reported infections were upper and lower respiratory tract infections. Grade 3-5 infections were reported in 11% of patients in the GAZYVA arm versus 10% of patients in the placebo arm. Fatal infection events were reported in 1% of patients in the GAZYVA arm versus 0.5% of patients in the placebo arm [see *Warnings and Precautions (5.6)*].

Thrombocytopenia

Chronic Lymphocytic Leukemia

The overall incidence of thrombocytopenia reported as an adverse reaction was higher in the GAZYVA treated arm (14%) compared to the rituximab product treated arm (7%), with the incidence of Grade 3 to 4 events being 10% and 3%, respectively (Table 7). The difference in incidences between the treatment arms is driven by events occurring during the first cycle. The incidence of thrombocytopenia (all grades) in the first cycle was 11% in the GAZYVA and 3% in the rituximab product treated arms, with Grade 3 to 4 rates being 8% and 2%, respectively. Four percent of patients in the GAZYVA treated arm experienced acute thrombocytopenia (occurring within 24 hours after the GAZYVA infusion).

The overall incidence of hemorrhagic events and the number of fatal hemorrhagic events were similar between the treatment arms, with 3 in the rituximab product and 4 in the GAZYVA treated arms. However, all fatal hemorrhagic events in patients treated with GAZYVA occurred in Cycle 1.

Non-Hodgkin Lymphoma

The incidence of thrombocytopenia in GADOLIN was lower in the GAZYVA plus bendamustine arm (15%) compared to the arm treated with bendamustine alone (25%). The incidence of hemorrhagic events in GAZYVA plus bendamustine treated patients compared to bendamustine alone was 12% and 11%, respectively. Grade 3 to 4 hemorrhagic events were similar in both treatment arms (4% in the GAZYVA plus bendamustine arm and 2% in the bendamustine arm).

In GALLIUM, thrombocytopenia was reported as an adverse reaction in 14% of the GAZYVA treated arm and 8% of the rituximab product treated arm, with the incidence of Grade 3 to 4 events being 7% and 3%, respectively. The difference in incidences between the treatment arms is driven by events occurring during the first cycle. The incidence of thrombocytopenia (all grades) in the first cycle was 9% in the GAZYVA and 3% in the rituximab product treated arms, with Grade 3 to 4 rates being 5% and 1%, respectively. In GALLIUM, both treatment arms had a 12% overall incidence of hemorrhagic events and a < 1% incidence of fatal hemorrhagic events.

Disseminated Intravascular Coagulation

In GALLIUM, DIC was reported as an adverse reaction in 0.3% of the GAZYVA treated patients. All events occurred within 1-2 days after the first infusion.

Tumor Lysis Syndrome

The incidence of Grade 3 or 4 tumor lysis syndrome in GAZYVA treated patients was 2% in CLL11, 0.5% in GADOLIN and 0.9% in GALLIUM.

Musculoskeletal Disorders

Chronic Lymphocytic Leukemia

Adverse reactions related to musculoskeletal disorders (all events from the body system), including pain, have been reported in the GAZYVA treated arm with higher incidence than in the rituximab product treated arm (18% vs. 15%).

Non-Hodgkin Lymphoma

In GADOLIN, adverse reactions related to musculoskeletal disorders (all events from the body system), including pain, have been reported in the GAZYVA plus bendamustine treated arm with higher incidence than in the bendamustine alone arm (44% vs. 30%).

In GALLIUM, musculoskeletal disorders were reported in 54% of patients in the GAZYVA treated arm and 49% of patients in the rituximab product treated arm.

Liver Enzyme Elevations

Hepatic enzyme elevations have occurred in CLL patients who received GAZYVA in clinical trials and had normal baseline hepatic enzyme levels (AST, ALT and ALP). The events occurred most frequently within 24–48 hours of the first infusion. In some patients, elevations in liver enzymes were observed concurrently with IRRs or tumor lysis syndrome. In the CLL11 study, there was no clinically meaningful difference in overall hepatotoxicity adverse reactions between all arms (4% of patients in the GAZYVA treated arm). Medications commonly used to prevent IRRs (e.g., acetaminophen) may also be implicated in these events. Monitor liver function tests during treatment, especially during the first cycle. Consider treatment interruption or discontinuation for hepatotoxicity.

Gastrointestinal Perforation

Cases of gastrointestinal perforation have been reported in patients receiving GAZYVA, mainly in NHL.

Worsening of Pre-existing Cardiac Conditions

Fatal cardiac events have been reported in patients treated with GAZYVA.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of GAZYVA. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- *Immune/Autoimmune Events*: Serum sickness

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings from animal studies and its mechanism of action, GAZYVA can cause fetal B-cell depletion [see *Clinical Pharmacology* (12.1)]. There are no data with GAZYVA use in pregnant women to inform a drug-associated risk. Monoclonal antibodies are transferred across the placenta. In animal reproduction studies, weekly intravenous administration of obinutuzumab to pregnant cynomolgus monkeys from day 20 of pregnancy until parturition which includes the period of organogenesis at doses with exposures up to 2.4 times the exposure at the clinical dose of 1,000 mg monthly produced opportunistic infections and immune complex mediated hypersensitivity reactions. No embryo-toxic or teratogenic effects were observed in the monkeys (*see Data*). Advise pregnant women of the potential risk to the fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown; however, the estimated background risk in the U.S. general population of major birth defects is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregnancies.

Clinical Considerations

Disease-Associated Maternal and/or Embryo/Fetal Risk

Pregnant women with systemic lupus erythematosus (SLE) are at increased risk of adverse pregnancy outcomes, including worsening of the underlying disease, premature birth, miscarriage, and intrauterine growth restriction. Maternal LN increases the risk of hypertension and preeclampsia/eclampsia. Passage of maternal autoantibodies across the placenta may result in adverse neonatal outcomes, including neonatal lupus and congenital heart block.

Fetal/Neonatal Adverse Reactions

GAZYVA is likely to cause fetal B-cell depletion (*see Data*). Avoid administering live vaccines to neonates and infants exposed to GAZYVA in utero until B-cell recovery occurs [*see Warnings and Precautions (5.11) and Clinical Pharmacology (12.2)*].

Data

Animal Data

In a pre- and post-natal development study, pregnant cynomolgus monkeys received weekly intravenous doses of 25 or 50 mg/kg obinutuzumab from day 20 of pregnancy until parturition, which includes the period of organogenesis. The high dose results in an exposure (AUC) that is 2.4 times the exposure in patients with CLL at the recommended label dose. There were no embryo-toxic or teratogenic effects in animals. Secondary opportunistic infections, immune complex mediated hypersensitivity reactions, or a combination of both were observed in exposed dams. When first measured on day 28 postpartum, obinutuzumab was detected in offspring at levels in the range of maternal serum levels on the same day, and B-cells were completely depleted. The B-cell counts returned to normal levels, and immunologic function was restored within 6 months after birth.

8.2 Lactation

Risk Summary

There is no information regarding the presence of GAZYVA in human milk, the effects on the breastfed child, or the effects on milk production. However, low levels of obinutuzumab were present in the milk of lactating cynomolgus monkeys [*see Data*]. Human IgG is known to be present in human milk. Because of the potential of serious adverse reactions in the breastfed child, advise women not to breastfeed during treatment with GAZYVA and for 6 months after the last dose.

Data

Obinutuzumab was measured in the milk of lactating cynomolgus monkeys on day 28 postpartum after weekly intravenous administration from day 20 of pregnancy until parturition. Concentrations in milk were approximately 0.04% and 0.13% of concentrations in maternal serum in the 25 and 50 mg/kg groups, respectively.

8.3 Females and Males of Reproductive Potential

GAZYVA can cause fetal harm when administered to a pregnant woman [*see Use in Specific Populations (8.1)*].

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment with GAZYVA and for 6 months after the last dose.

8.4 Pediatric Use

The safety and effectiveness of GAZYVA in pediatric patients have not been established.

8.5 Geriatric Use

Chronic Lymphocytic Leukemia

Of 336 patients with previously untreated CLL who received GAZYVA in combination with chlorambucil, 81% were 65 years and older, while 46% were 75 and older. Of the patients 75 years and older, 46% experienced serious adverse reactions and 7% experienced adverse reactions leading to death. Of the patients younger than 75, 33% experienced a serious adverse reaction and 2% an adverse reaction leading to death. No significant differences in efficacy were observed between younger and older patients [see *Clinical Studies (14.1)*].

Non-Hodgkin Lymphoma

Of 204 patients in GADOLIN with relapsed or refractory NHL treated with GAZYVA plus bendamustine, 44% were 65 and over, while 14% were 75 and over. In patients 65 and over, 55% of patients experienced serious adverse reactions and 28% experienced adverse reactions leading to treatment withdrawal while in patients under 65, 37% and 14% experienced serious adverse reactions and adverse reactions leading to treatment withdrawal, respectively. No clinically meaningful differences in efficacy were observed between these patients and younger patients in GADOLIN.

Of the 691 patients in GALLIUM treated with GAZYVA plus chemotherapy as first-line therapy, 33% were 65 and over, while 7% were 75 and over. Of patients 65 and over, 63% experienced serious adverse reactions and 26% experienced adverse reactions leading to treatment withdrawal, while in patients under 65, 43% experienced serious adverse reactions and 13% had an adverse reaction leading to treatment withdrawal. No clinically meaningful differences in efficacy were observed between these patients and younger patients in GALLIUM.

Lupus Nephritis

Clinical studies of GAZYVA in patients with active LN did not include sufficient numbers of patients aged 65 years and older (1 patient) to determine whether they respond differently from younger adult patients.

10 OVERDOSAGE

There has been no experience with overdose in human clinical trials. For patients who experience overdose, treatment should consist of immediate interruption or reduction of GAZYVA and supportive therapy.

11 DESCRIPTION

Obinutuzumab is a humanized anti-CD20 monoclonal antibody of the IgG1 subclass. It recognizes a specific epitope of the CD20 molecule found on B cells. The molecular mass of the antibody is approximately 150 kDa.

GAZYVA (obinutuzumab) injection is produced by mammalian cell (CHO) suspension culture. GAZYVA was engineered to have reduced fucose content as compared to a typical IgG1 produced in CHO cells. GAZYVA is a sterile, clear, colorless to slightly brown, preservative-free liquid concentrate for intravenous use. GAZYVA is supplied at a concentration of 25 mg/mL in 1,000 mg single-dose vials. Each vial contains in 40 mL: 1,000 mg obinutuzumab, L-histidine (57.6 mg), L-histidine hydrochloride monohydrate (89.6 mg), trehalose dihydrate (3632 mg), and poloxamer 188 (8 mg). The pH is 6.0.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Obinutuzumab is a monoclonal antibody that targets the CD20 antigen expressed on the surface of pre-B and mature B lymphocytes. Upon binding to CD20, obinutuzumab mediates B-cell lysis through (1) engagement of immune effector cells, (2) by directly activating intracellular death signaling pathways (direct cell death), and/or (3) activation of the complement cascade. The immune effector cell mechanisms include antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis.

As an antibody with reduced fucose content, obinutuzumab induces greater ADCC activity than rituximab in vitro using human cancer cell lines. Obinutuzumab also demonstrated an increased ability to induce direct cell death when compared to rituximab. Obinutuzumab binds to FcγRIII using purified proteins with a higher affinity than rituximab. Obinutuzumab and rituximab bind with similar affinity to overlapping epitopes on CD20.

12.2 Pharmacodynamics

In patients with CLL, GAZYVA caused CD19 B-cell depletion (defined as CD19 B cell counts $< 0.07 \times 10^9/L$). Initial CD19 B cell recovery was observed in some patients approximately 9 months after the last GAZYVA dose. At 18 months of follow-up, some patients remain B cell depleted.

Although the depletion of B cells in the peripheral blood is a measurable pharmacodynamic effect, it is not directly correlated with the depletion of B-cells in solid organs or in malignant deposits. B cell depletion has not been shown to be directly correlated to clinical response.

In patients with LN (REGENCY study), total peripheral CD19+ B cell levels below the defined threshold of 10 cells/ μl were achieved in 99% of patients treated with GAZYVA by Week 4 after treatment initiation and remained below this threshold in 95% of patients at Week 76.

Reductions in circulating naive B, memory B, and plasmablasts/plasma cells were observed by Week 4 and remained low through Week 76 after treatment initiation.

Treatment with GAZYVA led to improvements in complement (C3 and C4) and anti-double-stranded DNA antibodies (anti-dsDNA) by Week 4 and Week 12, respectively. These changes were sustained through Week 76.

In patients with low C3 at baseline, normalization of C3 levels occurred in 49% (by Week 12) and in 62% (by Week 76) of patients receiving GAZYVA compared to 33% and 29%, respectively, in the placebo group. In patients with low C4 at baseline, normalization of C4 levels occurred in 75% (by Week 12) and in 88% (by Week 76) of patients receiving GAZYVA compared to 55% and 55%, respectively, in the placebo group. Among patients with positive anti-dsDNA at baseline, 32% and 56% of patients treated with GAZYVA seroconverted by Week 4 and Week 76, respectively, compared with 16% and 16% of patients receiving placebo.

The clinical relevance of the above mentioned pharmacodynamic biomarkers has not been established.

Cardiac Electrophysiology

The potential effects of GAZYVA on the QTc interval have not been studied.

12.3 Pharmacokinetics

The pharmacokinetic parameters of obinutuzumab after 100 mg on day 1 and 900 mg on day 2 of Cycle 1, 1,000 mg on day 8 and 15 of Cycle 1, and 1,000 mg on day 1 of Cycles 2-6 for CLL; after 1,000 mg on day 1, 8 and 15 of Cycle 1, 1,000 mg on day 1 of Cycles 2-6 or Cycles 2-8, and then 1,000 mg every 2 months for up to 2 years for NHL; 1,000 mg on day 1, week 2, 24, 26 and every 6

months for up to 76 Weeks for LN are provided in Table 14. The dosing regimen is within the linear pharmacokinetic behavior of obinutuzumab.

Table 14 Obinutuzumab Pharmacokinetic Parameters

PK Measure	CLL ^a	Relapsed or refractory FL ^a	First line FL in combination with chemotherapy		LN ^c
			GAZYVA + Bendamustine ^a	GAZYVA + CHOP or CVP ^b	
C _{max} , µg/mL	466.3 (35)	553.5 (32)	513.4 (28)	676.4 (30)	463 (18)
C _{trough} , µg/mL	192.5 (78)	295 (56)	255 (46)	395 (44)	0.91 (752)
AUC, µg/mL*day	8701 (51)	11362 (41)	10088 (35)	10723 (37)	8770 (38)

Results are presented as geometric mean (% Coefficient of Variation).

^a Induction Cycle 6 of a 28-day cycle;

^b Induction Cycle 8 of a 21-day cycle.

^c Steady state values

Distribution and Elimination

CLL and NHL

The elimination of obinutuzumab is comprised of a linear clearance pathway and a time-dependent non-linear clearance pathway. As GAZYVA treatment progresses, the impact of the time-dependent pathway diminishes in a manner suggesting target-mediated drug disposition (TMDD) and saturation of the TMDD at the end of the treatment cycle at the proposed clinical dosing regimen. The distribution and elimination parameters of obinutuzumab in patients with CLL and NHL are provided in Table 15.

Table 15 Distribution and Elimination Parameters of Obinutuzumab

	CLL	NHL
Distribution		
Volume of Distribution ^a , L	4.1 (20)	4.3 (21)
Elimination		
Terminal Half-life, days	25.5 (48)	35.3 (35)
Clearance, L/day	0.11 (53)	0.08 (41)

Parameters are presented as geometric mean (% Coefficient of Variation).

^a At steady state.

LN

The steady state clearance of obinutuzumab was approximately 0.13 L/day with a median elimination t_{1/2} of 22.4 days.

GAZYVA elimination comprises two parallel pathways, a linear clearance pathway and a non-linear clearance pathway, which changes as a function of time. The time-varying clearance decreases with time with an exponential decay coefficient, likely related to CD20 target reduction and proteinuria improvement over time, and a time-independent clearance related to the endogenous catabolic processes of IgG.

Specific Populations

CLL and NHL

Age (median [range]: 63 [22, 89] years) and baseline creatinine clearance (CLcr) (median [range] 84 [22, > 120] mL/min) did not affect the pharmacokinetics of GAZYVA. In patients with CLcr $\leq$ 30 mL/min, the pharmacokinetics of GAZYVA was unaffected. GAZYVA has not been studied in patients with hepatic impairment.

The volume of distribution and steady-state clearance increased with body weight; however, the expected change in exposure does not warrant a dosage modification.

LN

The population pharmacokinetic analysis of GAZYVA showed that creatinine clearance (CLcr) (median [range] 108 [28.9 - 282 mL/min]) does not affect the pharmacokinetics in patients with LN. The pharmacokinetics of obinutuzumab in patients with mild or moderate renal impairment were similar to those in patients with normal kidney function. The safety and efficacy of GAZYVA in patients with severe renal impairment has not been formally studied.

12.6 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of anti-drug antibody (ADA) (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies or to other products may be misleading.

Seven percent (18/271) of patients with CLL tested positive for anti-GAZYVA antibodies at one or more time points in CLL11. No patients developed anti-GAZYVA antibodies during or following GAZYVA treatment in GADOLIN, while 1 patient (1/564, 0.2%) developed anti-GAZYVA antibodies in GALLIUM. Neutralizing activity of anti-GAZYVA antibodies was not assessed.

In GAZYVA-treated patients in the LN studies, a total of 12 out of 200 (6%) had at least one ADA-positive sample recorded at any time during the studies. Six (3%) subjects had ADA-positive samples recorded at baseline in which two patients remained ADA-positive throughout the studies, one patient had a single post-baseline sample that was ADA-positive, and three patients had post-baseline samples that were all ADA-negative. Six (3%) patients with ADA-negative samples at baseline had a positive ADA titer post-baseline (treatment-induced ADA). None of the 12 patients with positive ADA titers at any time during the treatment period experienced an IRR or hypersensitivity reaction during the studies. Neutralizing activity of anti-GAZYVA antibodies was not assessed.

Because of the low occurrence of anti-drug antibodies, the effect of these antibodies on the pharmacokinetics, pharmacodynamics, safety and/or effectiveness of GAZYVA is unknown.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No carcinogenicity or genotoxicity studies have been conducted with obinutuzumab.

No specific studies have been conducted to evaluate potential effects on fertility; however, no adverse effects on male or female reproductive organs were observed in the 26-week repeat-dose toxicity study in cynomolgus monkeys.

14 CLINICAL STUDIES

14.1 Chronic Lymphocytic Leukemia

The efficacy of GAZYVA was evaluated in a three-arm, open-label, active-controlled, randomized, multicenter trial (CLL11; NCT01010061) in 781 patients with previously untreated CD20+ CLL requiring treatment who had coexisting medical conditions or reduced renal function as measured by creatinine clearance (CLcr) < 70 mL/min. Patients with CLcr < 30 mL/min, active infections, positive hepatitis B (HBsAg or anti-HBc positive; patients positive for anti-HBc could be included if hepatitis B viral DNA was not detectable) and hepatitis C serology, or immunization with live virus vaccine within 28 days prior to randomization were excluded from the trial. Patients were treated with chlorambucil control (Arm 1), GAZYVA in combination with chlorambucil (Arm 2), or rituximab product in combination with chlorambucil (Arm 3). The safety and efficacy of GAZYVA was evaluated in a Stage 1 comparison of Arm 1 vs. Arm 2 in 356 patients and a Stage 2 comparison of Arm 2 vs. Arm 3 in 663 patients.

The majority of patients received 1,000 mg of GAZYVA on days 1, 8 and 15 of the first cycle, followed by treatment on the first day of 5 subsequent cycles (total of 6 cycles, 28 days each). The first dose of GAZYVA was divided between day 1 (100 mg) and day 2 (900 mg) [see *Dosage and Administration* (2.2)], which was implemented in 140 patients. Chlorambucil was given orally at 0.5 mg/kg on day 1 and day 15 of all treatment cycles (1 to 6).

In CLL11, the median age was 73 years, 62% were male, and 95% were White. Sixty-five percent had a CLcr < 70 mL/min and 76% had multiple coexisting medical conditions. Twenty-two percent of patients were Binet stage A, 42% were stage B, and 36% were stage C. The median estimated CLcr was 62 mL/min. Eighty-one percent of patients treated with GAZYVA in combination with chlorambucil received all 6 cycles compared to 89% of patients in the rituximab product treated arm and 67% in the chlorambucil alone arm.

In the Stage 1 analysis of CLL11, the median progression-free survival (PFS) in the GAZYVA in combination with chlorambucil arm was 27.2 months and 11.2 months in the chlorambucil alone arm (median observation time 22.8 months) as assessed by independent review and is consistent with investigator-assessed PFS. The median overall survival (OS) was not yet reached with a total of 46 deaths: 22 (9%) in the GAZYVA in combination with chlorambucil arm and 24 (20%) in the chlorambucil arm. The hazard ratio for OS was 0.41 (95% CI: 0.23-0.74).

In the Stage 2 analysis of CLL11, the median PFS was 26.7 months in the GAZYVA arm and 14.9 months in the rituximab product arm with a median observation time of 18.7 months (HR: 0.42, 95% CI: 0.33-0.54, p-value < 0.0001). These results were assessed by independent review and are consistent with investigator-assessed PFS. Minimal residual disease (MRD) was evaluated using allele-specific oligonucleotide polymerase chain reaction (ASO-PCR). The cutoff for a negative status was one CLL cell per 10⁴ leukocytes in the sample (i.e., an MRD value of < 10⁻⁴ was considered negative). Among patients who achieved complete response (CR) and complete response with incomplete marrow recovery (CRi; 94 patients in the GAZYVA arm and 34 patients in the rituximab product arm), 18 patients (19%) had negative MRD in the bone marrow in the GAZYVA arm compared to 2 patients (6%) in the rituximab product arm. Out of the patients who achieved CR and CRi, 39 patients (41%) in the GAZYVA arm, and 4 patients (12%) in the rituximab product arm

were MRD negative in peripheral blood samples collected at least 3 months after the end of treatment.

Efficacy results are shown in Table 16 and Figures 1 and 2.

Table 16 Efficacy Results from CLL11

Endpoint	Stage 1 of CLL11		Stage 2 of CLL11	
	GAZYVA + Chlorambucil* n = 238	Chlorambucil n = 118	GAZYVA + Chlorambucil* n = 333	Rituximab product + Chlorambucil n = 330
Median Progression-Free Survival ^a	27.2 months 11.2 months (HR 0.19 [0.14; 0.27], p-value < 0.0001 stratified log-rank test)		26.7 months 14.9 months (HR 0.42 [0.33; 0.54], p-value < 0.0001 stratified log-rank test)	
Overall Response Rate ^b	78.2%	33.1%	79.6%	66.3%
Complete Response	28.2%	0	26.1%	8.8%
Complete Response with Incomplete Marrow Recovery	2.5%	1.7%	2.1%	1.5%
Partial Response	45.0%	30.5%	48.6%	54.1%
Nodular Partial Response	2.5%	0.8%	2.7%	1.8%
Median Duration of Response	22.4 months	4.7 months	19.6 months	9.7 months
Overall Survival	HR 0.41 [0.23; 0.74]		Not Yet Mature	

^a As defined by independent review. Investigator-assessed PFS was consistent with data from independent review.

^b Defined as best overall response rate (ORR = CR + CRi + PR + nPR).

*All Stage 1 GCLb patients (n = 238) were included in the Stage 2 GCLb population (n = 333).

Figure 1

Kaplan-Meier Curve of Overall Survival in Patients with CLL in CLL11 (Stage 1)





Figure 2

Kaplan-Meier Curve of Progression-Free Survival in Patients with CLL in CLL11 (Stage 2)



14.2 Follicular Lymphoma

GADOLIN

The efficacy of GAZYVA was evaluated in GADOLIN (NCT01059630), an open-label, multicenter, randomized study that included 335 patients with follicular lymphoma (FL) who had no response to or have progressed during or within 6 months of rituximab product or a rituximab product-containing regimen. These patients were randomized to receive either bendamustine alone (n = 171) or GAZYVA in combination with bendamustine (n = 164) for 6 cycles, each of 28 days duration. Patients in the GAZYVA plus bendamustine arm who did not have disease progression [patients with a complete response (CR), partial response (PR) or stable disease (SD)] at the end of the 6 cycles continued receiving GAZYVA monotherapy for 2 years. Patients were stratified according to the type of refractoriness to rituximab product (refractory to rituximab product monotherapy versus rituximab product in combination with chemotherapy), the number of prior therapies (≤ 2 versus > 2), and geographic region.

GAZYVA was given by intravenous infusion as a flat dose of 1,000 mg on Days 1, 8 and 15 of Cycle 1, on Day 1 of Cycles 2–6, and then every 2 months until disease progression for up to 2 years. Bendamustine was given intravenously on Days 1 and 2 for all treatment cycles (1–6) at 90 mg/m²/day when given in combination with GAZYVA or 120 mg/m²/day when given alone.

The primary analysis included 321 FL patients, including 166 patients randomized to bendamustine alone and 155 patients randomized to GAZYVA in combination with bendamustine. In the primary analysis, patients had a median age of 63 years, 88% were White and 56% were male. Thirty-four percent had bulky disease (> 6 cm), 15% had at least one B-symptom at baseline and 95% had an ECOG performance status of 0–1 at baseline. The median time since initial diagnosis was 3 years and the median number of prior therapies was 2 (range 1 to 10). Forty-six percent of patients received 1 prior therapy and 33% of patients received 2 prior therapies. Twenty percent of patients were refractory to prior rituximab product monotherapy, 37% of patients were refractory to prior rituximab product plus chemotherapy induction treatment, and 41% of patients were refractory to rituximab product maintenance treatment received following rituximab product plus chemotherapy induction. Seventy-nine percent of patients were refractory to both rituximab product and an alkylating agent during any prior regimen (double refractory).

The major efficacy outcome measure was PFS as determined by an independent review committee (IRC). At the time of the primary analysis, median observation time was 21.1 months. The median PFS in the bendamustine arm was 13.8 months. Median PFS was not reached in the GAZYVA plus bendamustine arm (PFS HR = 0.48, 95% CI: 0.34-0.68; stratified log-rank test p-value < 0.0001). The investigator assessed PFS result was consistent with the IRC-assessed PFS. The median investigator-assessed PFS in the bendamustine arm was 13.7 months and the median in the GAZYVA containing arm was 29.2 months (PFS HR = 0.48, 95% CI: 0.35-0.67; stratified log-rank test p-value < 0.0001).

Efficacy results are summarized in Table 17. The Kaplan-Meier curve for IRC-PFS is shown in Figure 3.


Table 17 Primary Analysis Efficacy Results from GADOLIN^{a, b}

Endpoint	GADOLIN	
	GAZYVA + Bendamustine followed by GAZYVA monotherapy n = 155	Bendamustine n = 166
Median Progression-Free Survival (months)	Not Reached 13.8 (HR = 0.48 [0.34; 0.68], p-value < 0.0001 by stratified log-rank test)	
Best Overall Response ^c	78.7%	74.7%
Complete Response	15.5%	18.7%
Partial Response	63.2%	56.0%
Median duration of response (months)	Not Reached	11.6

^a Based on FL population.


^b As defined by independent review.

^c Best response of PR or CR within 12 months of study start.

Figure 3**Kaplan-Meier Curve of IRC-Assessed Progression-Free Survival in Patients with FL**

The final analysis included a total of 335 patients with 171 randomized to bendamustine alone and 164 to GAZYVA in combination with bendamustine. With an overall median observation time of 52.2 months (range: 0-100.9 months), there were 66 deaths (40.2%) in the GAZYVA arm and 85 deaths (51.3%) in the bendamustine-alone arm (OS HR = 0.71, 95% CI: 0.51, 0.98). The Kaplan-Meier curve for OS is presented in Figure 4.

Figure 4
Kaplan-Meier Curve of Overall Survival in Patients with FL



GALLIUM

The efficacy of GAZYVA was evaluated in GALLIUM (NCT01332968), a multicenter, open-label, randomized study that included 1202 patients with previously untreated, stage II bulky, III or IV FL. Patients were randomized 1:1 to receive either GAZYVA (n = 601) or rituximab product (n = 601) in combination with chemotherapy (CHOP, CVP, or bendamustine) for 6–8 cycles. Patients were stratified by chemotherapy (selected by each site; all patients at that site received the chosen chemotherapy regimen), FLIPI (Follicular Lymphoma International Prognostic Index) risk group and geographic region. Patients with at least PR to combination therapy received monotherapy with GAZYVA (1,000 mg) or rituximab product every two months until disease progression or for a maximum of two years. The study excluded patients with follicular lymphoma grade 3b or transformed disease; patients having an ANC < 1500 / μ L, platelets < 75,000 / μ L, or CLcr < 40 mL/min; and patients with hepatic transaminases > 2.5 x upper limit of normal unless attributable to lymphoma.

GAZYVA was given by intravenous infusion as a flat dose of 1,000 mg on Days 1, 8 and 15 of Cycle 1 and Day 1 of subsequent treatment cycles.

GAZYVA and bendamustine were given in six 28-day cycles. Bendamustine was administered at 90 mg/m²/day on Days 1 and 2 of each cycle, with prednisone 100 mg orally or equivalent on Day 1 of Cycle 1.

GAZYVA and CHOP were given in six 21-day cycles. Subsequently, two additional cycles of GAZYVA were given for a total of 8 GAZYVA cycles. CHOP consisted of cyclophosphamide 750 mg/m² intravenously, doxorubicin 50 mg/m², and vincristine 1.4 mg/m² (maximum dose, 2 mg) on Day 1 and prednisone 100 mg orally on Days 1-5.

GAZYVA and CVP were given in eight 21-day cycles. CVP consisted of cyclophosphamide 750 mg/m² intravenously and vincristine 1.4 mg/m² (maximum dose, 2 mg) on Day 1 and prednisone 100 mg orally on Days 1-5.

Patients had a median age of 59 years, 81% were White and 53% were female; 7% had Stage II, 35% had Stage III, and 56% had Stage IV disease, with 44% having bulky disease (≥ 7 cm) overall; 79% had a FLIPI score of > 2 ; and 97% had an ECOG performance status of 0–1. The chemotherapy was bendamustine in 57%, CHOP in 33%, and CVP in 10% of patients.

Efficacy was based on PFS per IRC, with a median observation time of 38 months. Upon interim analysis, the risk of progression or death was significantly reduced in the GAZYVA containing arm compared to the rituximab product containing arm (Table 18). Kaplan-Meier curves for PFS are shown in Figure 5. Overall response and complete remission rates were similar.

Table 18 Efficacy in Previously Untreated Follicular Lymphoma (GALLIUM)

Endpoint per IRC	GAZYVA + chemotherapy followed by GAZYVA monotherapy n = 601	Rituximab product + chemotherapy followed by rituximab product monotherapy n = 601
Progression-Free Survival ^a Number of events (%)	108 (18%) HR = 0.72 [95% CI: 0.56, 0.93], p-value = 0.0118 ^b	141 (23%)
Overall Response Rate ^c	91%	88%
Complete Remission Rate ^c	28%	27%


^a Investigator-assessed PFS was consistent with data from independent review.

^b Stratified log-rank test.

^c After completion of combination therapy. Assessed by CT without positron emission tomography.

Figure 5

Kaplan-Meier Curves of Progression Free Survival in Patients with Previously Untreated FL



CI, confidence interval; G-chemo, obinutuzumab plus chemotherapy; HR, hazard ratio; NR, not reached; PFS, progression-free survival; R-chemo, rituximab plus chemotherapy

14.3 Active Lupus Nephritis

Study Design and Population

The efficacy of GAZYVA was evaluated in REGENCY (NCT04221477), a Phase III, randomized, double-blind, placebo-controlled, parallel-group, multicenter study in 271 patients with ISN/RPS 2003 Class III or IV, with or without concomitant Class V lupus nephritis (LN), treated with standard therapy consisting of mycophenolate mofetil (MMF) and corticosteroids. Patients had active or active/chronic ISN/RPS 2003 Class III or IV, with or without concomitant Class V proliferative LN determined by kidney biopsy, current or past positive antinuclear antibody (ANA), urine protein-to-creatinine ratio (UPCR) ≥ 1 g/g, and had received at least one dose of pulse intravenous (IV) methylprednisolone (≥ 250 mg) or equivalent treatment for LN during the 6 months prior to screening or during screening.

Patients with estimated glomerular filtration rate (eGFR) <30 ml/min/1.73 m² or in need of dialysis or transplantation, with sclerosis in $> 50\%$ of glomeruli on kidney biopsy, presence of rapidly progressive glomerulonephritis, evidence of active infection, receipt of anti-CD20 therapy < 9 months before or during screening, or receipt of cyclophosphamide, tacrolimus, ciclosporin, or voclosporin within 2 months of or during screening were excluded.

Patients were randomized 1:1 to receive GAZYVA 1,000 mg (N=135) or placebo (N=136) intravenously, in combination with MMF 2-2.5 g/day and a tapering course of corticosteroids and were evaluated over 76 weeks. Patients randomized to receive GAZYVA were further randomized in a 1:1 ratio to receive either GAZYVA 1,000 mg IV on Day 1, Weeks 2, 24, 26, 50, and 52 (Arm 1), or GAZYVA 1,000 mg IV on Day 1, Weeks 2, 24, 26, and 52 (Arm 2). The totality of the GAZYVA efficacy data combining both treatment arms is shown in Table 19.

All patients received oral prednisone 0.5 mg/kg/day (maximum 60 mg/day) and remained at this dose until Week 2. Beginning on Day 15, prednisone was tapered to achieve a target dose of 5 mg/day by Week 24. Prednisone was maintained at a low dose (5 mg/day) from Week 24 until Week 80.

The median age of patients was 31 years, 85% were female, 58% were Hispanic or Latino, 48% were White, 19% were American Indian or Alaska Native, 15% were Black or African American and 6% were Asian. The distribution by kidney biopsy class was 39% Class III, 61% Class IV and 31% had concomitant Class V. Mean (SD) eGFR at baseline 102.3 (± 30.8) mL/min/1.73 m². Mean (SD) UPCR at baseline was 3.3 (± 2.9) mg/mg with 42% of patients exhibiting UPCR ≥ 3 mg/mg at baseline.

Efficacy Results

The primary endpoint measure was proportion of patients who achieved complete renal response (CRR) at Week 76, defined as meeting all of the following criteria: UPCR < 0.5 g/g; eGFR $\geq 85\%$ of baseline, as calculated using the 2009 Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation; with no occurrence of the following intercurrent events: rescue therapy, treatment failure, death or early study withdrawal.

Key secondary endpoint measures included: proportion of patients who achieved CRR with successful prednisone taper at Week 76 (defined as achievement of CRR at Week 76 without receiving prednisone > 7.5 mg/day or equivalent from Week 64 through Week 76), proportion of patients who achieved a proteinuric response at Week 76 (defined as achievement of UPCR < 0.8 g/g and no occurrence of the following intercurrent events: rescue therapy, treatment failure, death or early study withdrawal) and proportion of patients who experience “renal-related events or deaths” through Week 76 (defined as occurrence of death, treatment failure, $\geq 50\%$ increase in UPCR to a value ≥ 3 g/g and/or $\geq 30\%$ decrease in eGFR to < 60 ml/min/1.73 m²).

The proportion of patients achieving CRR at Week 76 was significantly greater in patients treated with GAZYVA in combination with standard therapy compared to patients who received placebo plus standard therapy. There were also a higher proportion of patients who achieved CRR with successful prednisone taper at Week 76 and proteinuric response at Week 76 in the GAZYVA plus standard therapy arm compared to the placebo plus standard therapy arm (see Table 19).

In the REGENCY study, patients who received GAZYVA were less likely to experience the outcome of “renal-related event or death” compared with placebo. Fewer patients in the GAZYVA arm experienced worsening kidney function or doubling of serum creatinine (see Table 20).

Table 19 Summary of Efficacy Results in Adult Patients with Active Lupus Nephritis (REGENCY Study)

	GAZYVA + standard therapy (N=135)	Placebo + standard therapy (N=136)
Primary Endpoint		
<i>Complete renal response (CRR) at Week 76 (%)</i>	46.4 (38.0, 54.9)	33.1 (25.2, 41)
Treatment difference (95% CI) ^a	13.4 (2.0, 24.8)	
p-value	0.0232	
Components of CRR:		
UPCR < 0.5 g/g	64 (47.4%)	49 (36.0%)
eGFR ≥ 85% at baseline	113 (83.7%)	103 (75.7%)
No occurrence of intercurrent events	120 (88.9%)	102 (75%)
Key Secondary Endpoints		
<i>CRR with successful prednisone taper at Week 76 (%)</i>	42.7 (34.3, 51.1)	30.9 (23.1, 38.7)
Treatment difference (95% CI)	11.9 (0.6, 23.2)	
p-value	0.0421	
<i>Proteinuric response at Week 76 (%)</i>	55.5 (47.1, 64)	41.9 (33.6, 50.2)
Treatment difference (95% CI)	13.7 (2.0, 25.4),	
p-value	0.0227	

^a Primary and key secondary endpoints were analyzed using the Cochran-Mantel-Haenszel (CMH) test, adjusted for stratification factors race and region. Multiple Imputation handled missing data. For the primary endpoint CRR and the key secondary endpoint CRR with successful prednisone taper, missing data (not due to an Intercurrent Event (ICE)) occurred in four patients in the GAZYVA arm and one in the placebo arm. For the key secondary endpoint proteinuric response, four GAZYVA patients and two placebo patients had non-ICE-related missing data.

Table 20 Renal-Related Event or Death in Adult Patients with Active Lupus Nephritis (REGENCY Study) by Week 76

	Week 0-76		Hazard Ratio (HR) vs. Placebo (95% CI) Week 76
	GAZYVA + standard therapy N=135 (%)	Placebo + standard therapy N=136 (%)	
Renal-Related Event or Death			
Number (%) of patients with event	24 (17.8%)	46 (33.8%)	
Time to event			0.5 (0.3, 0.8)
Components of Renal-Related Event or Death Endpoint			
Percentage of patients with:			
Death	3 (2.2%)	1 (0.7%)	
Treatment failure ^a	6 (4.4%)	25 (18.4%)	
End-stage renal disease (ESRD) ¹	0	2 (1.5%)	
Clinically significant, sustained worsening in UPCR and/or eGFR from Week 24 ²	5 (3.7%)	22(16.2%)	
Rescue Therapy except corticosteroid-only rescue ³	5 (3.7%)	22 (16.2%)	
Worsening proteinuria ^b	17 (12.6%)	18 (13.2%)	
Worsening eGFR ^c	7 (5.2%)	20 (14.7%)	
Additional Renal-Related Events			
Percentage of patients with event			
Doubling of serum creatinine from baseline through Week 76	4 (3.0%)	8 (5.9%)	

^a Treatment failure was prospectively defined as any of the following: 1) new ESRD or need for chronic dialysis or renal transplantation, 2) clinically significant, sustained worsening in UPCR and/or eGFR from Week 24 onward that leads the investigator to conclude the patient failed the randomized treatment period, or 3) receipt of rescue therapy, except corticosteroid-only rescue


^b Worsening proteinuria was prospectively defined as a confirmed $\geq 50\%$ increase in UPCR to a value ≥ 3 g/g

^c Worsening eGFR was prospectively defined as confirmed $\geq 30\%$ decrease in eGFR to a value < 60 mL/min/1.73m²

Subgroup Analyses

In pre-specified subgroup efficacy analyses, the primary endpoint in patients was examined based on 24-hour UPCR at baseline (< 3 g/g or ≥ 3 g/g), lupus nephritis class at baseline (Class III or IV), and concomitant Class V at baseline. The results are displayed in Figure 6 below.

Figure 6
Forest Plot of CRR at Week 76 on Baseline Disease Characteristics, Efficacy-Evaluable Patients



Subgroup findings should be interpreted with caution due to small sample sizes and overlapping subgroups.
All patients received standard of care, consisting of MMF and corticosteroids, as per protocol.

16 HOW SUPPLIED/STORAGE AND HANDLING

GAZYVA (obinutuzumab) injection is a clear, colorless to slightly brown, preservative-free solution for intravenous use supplied as 1,000 mg/40 mL (25 mg/mL) in single-dose vials (NDC 50242-070-01).

Store at 2°C to 8°C (36°F to 46°F). Do not use beyond expiration date stamped on carton. Protect from light. DO NOT FREEZE. DO NOT SHAKE.

17 PATIENT COUNSELING INFORMATION

Infusion-related Reactions

Advise patients to seek immediate medical attention if signs and symptoms of infusion-related reactions occur [see *Warnings and Precautions* (5.3) and *Adverse Reactions* (6.1 and 6.2)].

Tumor Lysis Syndrome

Advise patients to seek immediate medical attention if symptoms of tumor lysis syndrome occur [see *Warnings and Precautions* (5.5) and *Adverse Reactions* (6.1)].

Infections

Advise patients to seek immediate medical attention if signs of an infection occur [see *Warnings and Precautions* (5.6) and *Adverse Reactions* (6.1 and 6.2)].

Hepatitis B Virus Reactivation

Advise patients to seek immediate medical attention if symptoms of hepatitis occur [see *Warnings and Precautions* (5.1)]. Advise patients of the need for hepatitis B virus screening and monitoring during treatment with GAZYVA.

Progressive Multifocal Leukoencephalopathy

Advise patients to seek immediate medical attention if new or changes in neurological symptoms occur [see *Warnings and Precautions* (5.2)].

Neutropenia, Thrombocytopenia and Disseminated Intravascular Coagulation

Advise patients to seek immediate medical attention if signs and symptoms of bleeding or thrombosis occur. Advise patients of the need for periodic monitoring of blood counts [see *Warnings and Precautions* (5.7, 5.8 and 5.9) and *Adverse Reactions* (6.1 and 6.2)].

Immunization

Advise patients to avoid vaccinations with live viral vaccines [see *Warnings and Precautions* (5.10)].

Embryo-Fetal Toxicity

Advise pregnant women of potential fetal B-cell depletion. Advise females of reproductive potential to inform their healthcare provider of a known or suspected pregnancy [see *Warnings and Precautions* (5.11), *Use in Specific Populations* (8.1)].

Contraception

Advise females of reproductive potential to use effective contraception during treatment with GAZYVA and for 6 months after the last dose [see *Use in Specific Populations* (8.3)].

Lactation

Advise women not to breastfeed during treatment with GAZYVA and for 6 months after the last dose [see *Use in Specific Populations* (8.2)].

GAZYVA® (obinutuzumab)

Manufactured by:

Genentech, Inc.

A Member of the Roche Group

1 DNA Way

South San Francisco, CA 94080-4990

U.S. License No. 1048

GAZYVA is a registered trademark of Genentech, Inc.

© 20XX Genentech, Inc.